

ON THE EVIDENCE FOR COSMIC VARIATION OF THE FINE STRUCTURE CONSTANT: A BAYESIAN REANALYSIS OF THE QUASAR DATASET

BY EWAN CAMERON* AND TONY PETTITT

Queensland University of Technology

We review the evidence behind recent claims of spatial variation in the fine structure constant deriving from observations on ground-based telescopes of ionic absorption lines in the light from distant quasars. To this end we expand upon previous non-Bayesian analyses limited by the assumptions of a strictly Normal and unbiased form for the “unexplained errors” of the benchmark quasar dataset. Through nested importance sampling and the method of power posteriors we evaluate and compare marginal likelihoods (or Bayes factors) for three competing hypotheses—(I) the strict null (no cosmic variation), (II) the monopole null (a constant Earth-to-quasar offset only), and (III) the monopole+dipole hypothesis (featuring a cosmic variation manifest to the Earth-bound observer as a North–South divergence)—under various alternative error terms. Our analysis reveals significant support for a skeptical interpretation in which the apparent dipole effect is driven solely by systematic errors of opposing sign inherent in measurements from the two telescopes employed to obtain the observations. In this context we highlight the importance of new observations along the equator of the alleged dipole (in addition to the new polar observations planned) in order to more strongly test the skeptical interpretation.

1. Introduction. Introduced in 1916 by Arnold Sommerfeld to abbreviate a recurring, dimensionless factor in his relativistic extension of the Bohr model for the hydrogen atom the fine structure constant, α , is now recognized as one of the fundamental coupling terms of quantum electrodynamics (QED). In this context α serves to characterize the strength of the electromagnetic interaction and has been humorously dubbed, “the Peter Sellers of QED” (Borie, 1986), owing to the many different roles it plays within this theory: setting the scale for all electromagnetic cross-sections, binding energies, and decay rates; and, of course, the scale of fine structure splitting in atomic and ionic spectral lines (cf. Griffiths 2005).

Defined by the ratio of the squared elementary charge, e^2 , to the permittivity of free space, ϵ_0 , the reduced Planck constant, $\hbar$, and the vacuum

*E.C. is grateful for financial support from the Australian Research Council.

Keywords and phrases: Bayesian inference, Cosmology

speed of light, c ,

$$(1) \quad \alpha = \frac{e^2}{(4\pi\epsilon_0)\hbar c},$$

the fine structure constant has an on-Earth laboratory value, known to remarkable precision through exacting experimentation on quantum scale systems, of $\alpha^{-1} \approx 137.035999037(91)$ (Bouchendira et al., 2011; Hanneke, Fogwell and Gabrielse, 2008). Though nominally designated a “constant” there are in fact substantial theoretical foundations to support a time-/space-varying α , if empirical evidence of such can be definitively established; most notably, within electrodynamic scalar field models (Bekenstein, 1982; Carroll, 1998) and grand unification theories (Marciano, 1984; Brax et al., 2003). An historical evolution (with respect to cosmological time) of the fine structure constant driven by a decreasing speed of light could even offer a compelling solution to the infamous “Horizon Problem”¹ of Big Bang cosmology without inflation (Moffat, 1993; Barrow, 1999; Albrecht and Magueijo, 1999).

Intriguingly, there exist a number of different approaches, both terrestrial and astronomical, through which one may search for this experimentalist’s “Holy Grail”. These include: (I) the in-laboratory comparison of optical atomic clocks (Fortier et al., 2007; Rosenband et al., 2008); (II) the nuclear modeling of samarium isotopes extracted from the Oklo natural fission reactor (Shlyakhter, 1976; Damour and Dyson, 1996; Gould et al., 2006) or rhenium extracted from Earth-fallen meteorite samples (Olive et al., 2004); (III) the cosmological modeling of angular fluctuations in the temperature and polarization of the Cosmic Microwave Background (Rocha et al., 2004; Nakashima, Nagata and Yokoyama, 2008; Calabrese et al., 2011); and/or (IV) the search for frequency shifts in α -sensitive features of astronomical spectra, most notably the fine structure emission lines of ionized carbon observed in radio waves from distant lensed galaxies (Levshakov et al., 2012; Weiss et al., 2012) and the ionic absorption lines of various species observed in the visible/near-visible light from distant quasars (Webb et al., 1999, 2001; Murphy et al., 2001; Murphy, Webb and Flambaum, 2003; Webb et al., 2011; King et al., 2012). After a number of initial disagreements between rival teams—see Lamoreaux and Torgerson (2004), for instance—now resolved, only the post-Millennial quasar-based studies at present claim to have recov-

¹Namely, the remarkable homogeneity of the Universe beyond even the scale of casual connection under the standard model; that is, over physical separations exceeding the maximum distance traversable since the beginning of time at the present-day speed of light.

ered any significant evidence for an evolving fine structure constant. However, the Webb et al. team’s interpretation of their quasar dataset in this regard remains a matter of ongoing controversy; the statistical analysis of which is the subject of this paper.

First identified in 1966 (Burbidge, Lynds and Burbidge, 1966; Stockton and Lynds, 1966) the distinctive absorption lines apparent to Earth-bound observers at UV-to-optical wavelengths in the light from distant quasars (or “quasi-stellar radio sources”) arise from the interaction of this light with ions of various species encountered during its passage through intervening gas clouds; these residing in the extensive halos of both bright, star-forming galaxies and dark proto-galaxies. The potential of these absorption lines (especially the singly and triply ionized silicon [SiII and SiIV] doublets) as probes of the fundamental constants at extra-galactic distances, complementary to the already-used emission line technique (Savedoff, 1956; Bahcall and Salpeter, 1965), was quickly realized by Bahcall, Sargent and Schmidt (1967), and used to constrain the cosmic variation of α to within $\sim 10\%$ of its on-Earth value ($\Delta\alpha/\alpha = 0.98[0.05]$) in the direction of one particular quasar (3C 191). (Throughout both the astronomical literature and our analysis herein the particular notation, $\Delta\alpha/\alpha$, is used to represent the fractional offset of the fine structure constant in the extra-galactic system under study from its on-Earth laboratory value, i.e., $\Delta\alpha/\alpha = [\alpha - \alpha_{\text{Earth}}]/\alpha_{\text{Earth}}$.) Improvements in astronomical instrumentation have since allowed far stronger constraints on α variation to be established through this approach. Ivanchik, Potekhin and Varshalovich (1999), for instance, have demonstrated $|\Delta\alpha/\alpha| < 2.3 \times 10^{-4}$ (95% CI) in the early Universe (at ~ 4 Gyr after the Big Bang; ~ 9.4 Gyr ago) through SiIV doublet observations along nine quasar sightlines from a ground-based Russian telescope (the BTA-6 at the Special Astrophysical Observatory).

To search for cosmic α variation below this limit (of roughly one part in ten thousand) with quasar absorption line spectra requires application of the “many multiplet” (MM) technique (Dzuba, Flambaum and Webb, 1999) in which the relative frequency shifts of *multiple* ionic species are simultaneously compared; the transitions of those species relatively insensitive to α variation (e.g. singly ionized magnesium [MgII]) effectively serving as calibration benchmarks for the stronger shifters (e.g. singly ionized iron [FeII]). Implementation of the MM technique necessarily involves a multi-parameter fit to constrain not only $\Delta\alpha/\alpha$ but also a suite of nuisance parameters accounting for the physical structure of the intervening gas cloud (i.e., column density, kinetic temperature, and the dominant line broadening mechanism). Pioneering this technique in 1999 the Webb et al. team (Webb et al., 1999)

recovered *tentative evidence that α may have been lower in the past* from a sample of 30 extra-galactic absorbers spanning $0.5 < z_{\text{abs}} < 1.6$ (here z_{abs} denotes the cosmological redshift² of the absorber(s) under study) observed with the Keck telescope ($\Delta\alpha/\alpha = -1.1 [0.4] \times 10^{-5}$). This exciting result was quickly heralded as support for a certain class of cosmological model admitting a decelerating speed of light (Barrow and Magueijo, 2000; Davies, Davis and Lineweaver, 2002), though the claimed theoretical basis for the well-publicized Davies et al. interpretation—derived from a (mistaken) consideration of black hole thermodynamics—was soon disproven (Carlip and Vaidya, 2003). More interestingly from a statistical point of view was the Webb et al. team’s concern for a possible underestimation of the true errors in their quoted uncertainties, with a remarkably strong dip in their α estimates over a narrow redshift interval ($0.9 \lesssim z_{\text{abs}} \lesssim 1.1$) suggesting the presence of an unexplained source of observational error.

Further post-Millennial studies by the same team (Webb et al., 2001; Murphy et al., 2001; Murphy, Webb and Flambaum, 2003) with the Keck telescope—expanding their original sample to a total of 141 absorbers and their redshift baseline to $0.2 < z_{\text{abs}} < 3.7$ —ultimately strengthened the apparent weight of evidence for a time-varying fine structure constant beyond the “ 4σ level”; given the particular assumptions of the statistical analysis employed. Chief among these the absence of any *bias* in the afore-mentioned unexplained error term—its presence still inferred from a marked excess of the sample variance over that expected from known sources of observational noise, and seemingly greater among the high redshift population. Murphy, Webb and Flambaum (2003) thus directed their attention at this time toward a number of possible explanatory factors for an additional error term unique to high redshift systems—including the entry of damped Lyman-alpha absorbers (dense clouds of neutral hydrogen featuring complex velocity structures more challenging to model via the MM technique) into the sample at $z_{\text{abs}} \gtrsim 2$ (where the rest-frame ultra-violet of their characteristic spectral lines is redshifted within the optical window accessible to on-Earth

²The cosmological redshift—literally a frequency shift “redwards” (toward longer wavelengths) apparent to Earth-based observers in the light received from extra-galactic systems due to cosmological expansion of the intervening space—serves as a proxy for lookback distance (or lookback time) in astronomy: the greater an object’s observed redshift the greater the physical separation between us and it at the time the light received on Earth was emitted, and the longer this light has had to travel to reach us. Integral formulae relating these physical quantities (subject to a specified parameterization of the standard cosmological model) are provided by Hogg (1999). For reference, we note that $z = 0$ corresponds (of course) to the present day, $z = 1$ to a lookback distance, $r(z)$, of ≈ 7.8 GLyr, and $z = 3$ to $r(z) \approx 11.4$ GLyr.

observers). Yet another, the imprint of a spatial variation in α (as we describe below), was discounted at this stage as only weakly supported by the available data (according to a bootstrap significance test); see [Murphy, Webb and Flambaum \(2003\)](#).

It was thus a great surprise in 2010 when new $\Delta\alpha/\alpha$ estimates for 154 (primarily southern hemisphere) absorption systems along 60 quasar sight-lines (52 new and 8 in common with the original Keck sample) observed by the same team ([King et al., 2012](#); [Webb et al., 2011](#))—but this time using the Very Large Telescope (VLT) in Chile—appeared to indicate the opposite evolutionary trend with redshift. Namely, *that the fine structure constant was in fact higher in the past for these absorbers* ($\Delta\alpha/\alpha = 0.154 [0.132] \times 10^{-5}$). To resolve this contradiction the team were forced to resurrect the spatial variation hypothesis, proposing a smooth transition in α across the Universe manifest to the on-Earth observer as a (z -invariant) monopole plus a (z -dependent³) dipole field. We illustrate graphically in Figure 1 (over page) the nature of the spatial variation in $\Delta\alpha/\alpha$ under the best-fit model of this form (cf. Section 2.3) from [King et al. \(2012\)](#) on a color-/symbol-coded map of the celestial sphere (represented here via a Mollweide projection in the equatorial coordinate system of right ascension and declination⁴). Note the scale of the inferred variation, which is at the $\Delta\alpha/\alpha \lesssim 10^{-5}$ level only

³[King et al. \(2012\)](#) in fact consider a variety of candidate functional forms for their dipole model, including a (redshift) z -invariant dipole, a z^β -dependent dipole, and an $r(z)$ -dependent dipole (with $r(z)$ the cosmological lookback distance). All exhibit (bootstrap randomization-based) statistical significances of $\sim 4\sigma$ over a monopole-only null. However, we note that: (i) a z -invariant dipole strictly implies a breaking of the Copernican principle—namely, that the Earthian observer does not occupy a privileged position within the cosmos—in addition to the cosmological principle—namely, that the physical laws of the Universe are uniform throughout; and consequently that the Universe appears essentially homogenous when viewed on sufficiently large scales—broken by all; and (ii) of the other two, the z^β - alternative (which encompasses a close approximation to the $r(z)$ -model at $\beta \approx 0.3$) already appears from the [King et al. \(2012\)](#) study to be an over-fitting of the available data. Hence, we focus exclusively on the (monopole+) $r(z)$ -dipole scenario in the present analysis. For reference, this was also the approach taken by [Berengut, Kava and Flambaum \(2012\)](#).

⁴The equatorial coordinate system charts the position of astronomical bodies on the celestial sphere with respect to the Earth’s equatorial plane. Right ascension (typically quoted in units of “time” from 0 to 24 hours) measures the angular distance *along the equator* from the vernal equinox to the (uniquely-specified) great circle passing through the designated object and the celestial poles, while declination (typically quoted in degrees from -90° to 90°) measures the corresponding angular distance *along this great circle* from the equator. Due to the slow, but astronomically appreciable, precession of the Earth’s orbital axis it is necessary to define an epoch of reference for this coordinate system; the present convention is to quote all equatorial coordinates as if observed at noon on 1 January 2000 (denoted the J2000 epoch).

accessible (as noted earlier) via the MM technique.




FIG 1. Visualization of the monopole+ $r(z)$ -dipole model (cf. Section 2.3) for explaining the apparent spatial variation of the fine structure constant proposed by the Webb et al. team. The fractional difference of the fine structure constant from its on-Earth value under this model, $\Delta\alpha/\alpha_{\text{mod}}$, as a function of the observational sightline at $z = 1$ under the best-fit solution of King et al. (2012) is shown in the lefthand panel and that at $z = 3$ in the righthand panel (with the color-/symbol-coding explained in the top-right legend of each). The maximum, minimum, and equator of the key North–South dipole component of this model are overlaid as well for reference.

In Figure 2 (over page) we present a comparable visualization of the “raw” $\Delta\alpha/\alpha$ variation in the Webb et al. team’s *quasar dataset* from which the apparent dipole effect shown above was inferred. To this end we compute in each bin of right ascension and declination containing at least one quasar sightline (but typically two or more; noting as well that there are on average 2–3 absorbers per sightline) the weighted mean,

$$\overline{\Delta\alpha/\alpha} = \frac{\sum (\Delta\alpha/\alpha) / (\sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2)}{\sum 1 / (\sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2)}.$$

Here σ_{ran} represents the standard deviation of the unexplained error term in each of the three ε_{ran} groups under the Webb et al. team’s proposed generative model (which we detail in Section 2.2). Despite the substantial degree of noise evident in these measurements (note the increase in scale with respect to that of Figure 1) one may yet discern an excess of negative $\Delta\alpha/\alpha$ estimates in the far North and an excess of positive $\Delta\alpha/\alpha$ estimates in the far South as per the dipole hypothesis. (Though, in anticipation of the skeptical interpretation of these results [i.e., that biases of opposite sign in the measurements from each telescope are to blame for the apparent North–South divergence], we note also that the equator of the alleged dipole

provides a near perfect subdivision of the sample into its VLT and Keck constituents.)




FIG 2. Visualization of the apparent dipole signature in the [King et al. \(2012\)](#) / [Webb et al. \(2011\)](#) quasar dataset. Each quasar sightline in the sample is marked here on an equal-area Mollweide projection of the sky based on the J2000 equatorial coordinate system; with solid-symbols denoting VLT observations and open symbols Keck observations. Note that for most of these quasars there are multiple intervening absorbers providing independent measurements of $\Delta\alpha/\alpha$ along the sightline. The apparent spatial variation of this observable is illustrated via a color-/symbol-coding (detailed in the top-right legend) of its weighted mean in each of the indicated subdivisions. The maximum, minimum, and equator corresponding to the best-fit monopole+ $r(z)$ -dipole solution of the [King et al. \(2012\)](#) paper are overlaid as well for reference.

The intriguing hypothesis of a spatial variation in the quasar dataset with monopole+ $r(z)$ -dipole form has been identified by [King et al. \(2012\)](#) through a bootstrap randomization (of $\Delta\alpha/\alpha$ estimates across sightlines and surveys) as favored over the simple monopole (i.e., time-/space-invariant with constant [non-zero] Earth-to-quasar offset) alternative at the “ 4σ significance level”—a result echoed by [Berengut, Kava and Flambaum \(2012\)](#) in their review of the dataset using the Akaike Information Criterion, the F statistic,

and the “error ellipsoid method”. With the new VLT sample apparently just as subject to unexplained measurement noise as the original Keck dataset, however, a number of strong assumptions were required to justify the significance testing procedures employed in this work. In particular, the authors were obliged to treat the aforesaid as strictly Normal and *strictly unbiased* (*viz.* zero mean and mode). Assessing the impact of these assumptions, which are easily relaxed within a Bayesian framework as we demonstrate herein, thus represents an essential “next step” in the analysis of this observational benchmark given the profound implications for both fundamental physics and cosmology if the dipole interpretation is to become accepted.

Following the extensive world-wide media coverage of their work⁵ the Webb et al. team’s spatial variation hypothesis was strongly criticized in a number of public forums, including the popular science blogs, “Uncertain Principles” by [Chad Orzel](#) and “Cosmic Variance” by [Sean Carroll](#). The former citing the evident alignment of the alleged dipole’s equator with the coverage overlap of the Keck and VLT telescopes (cf. Figure 2) as an indicator that biases of opposite sign in the observations from each might well be to blame; and the latter citing the extraordinarily low mass-energy budget ($\sim 10^{-42}$ GeV) for the underlying scalar field implied by the cosmic expanse of the fitted dipole. From a Bayesian perspective these objections may be viewed as disagreements over the relative prior probabilities of competing proposals. Namely, the degree to which the particular model for the unexplained error term adopted by the experimenters should be favored over some alternative model (or family of models), and the degree to which the null hypothesis (of a time-/space-invariant α) should be favored “theoretically” over the proposed dipole hypothesis.

In this paper we aim to address the above concerns explicitly within the Bayesian framework by computing, and thereby comparing, the marginal likelihoods of the dipole hypothesis, the monopole-only null hypothesis, and the strict null hypothesis (of $\Delta\alpha/\alpha$ everywhere zero) under both the Webb et al. team’s nominal error model and a selection of more skeptical alternatives. In Section 2 we summarize the contents of the quasar dataset, discuss the *explained* and *unexplained* sources of uncertainty in these measurements, and detail the functional form of the proposed monopole+ $r(z)$ -dipole. We then motivate our exploration of non-Normal and biased error terms through an inspection of the residuals about the best-fit dipole solution from [King et al. \(2012\)](#) and a comparison of $\Delta\alpha/\alpha$ estimates for “twice-observed” absorbers. In Section 3 we propose data-driven error models for the quasar dataset

⁵See, for example, the 23 October 2010 issue of New Scientist magazine (#2783), or the 2 September 2010 issue of The Economist magazine.

and specify priors on their controlling parameters. We then specify priors on the fundamental parameters of the competing hypotheses, and explore the resulting posterior probability densities through Markov Chain Monte Carlo analysis. In Section 4 we compute marginal likelihoods for each model via nested importance sampling and the method of power posteriors, and examine the sensitivity of the corresponding Bayes factors to our priors. Finally, in Section 5 we discuss the implications of these results in terms of the evidence, or lack thereof, for a spatial variation in the fine structure constant, and the prospects for future observations to settle this debate.

2. The Quasar Dataset. In this study we examine the publicly-available quasar dataset of Webb et al. (2011): a single catalog listing for each of the 295 intervening absorbers (141 Keck plus 154 VLT) identified along the 131 quasar sightlines probed to-date their MM-based estimates of z_{abs} , $\Delta\alpha/\alpha$, and σ_{obs} ; the last a standard deviation characterizing the uncertainty in each $\Delta\alpha/\alpha$ estimate arising from *explained* sources of observational error. Also listed are the J2000 identifier (encoding right ascension and declination in the equatorial coordinate system) and redshift, z_{qua} , of each background quasar, the telescope on which the corresponding observation was made, and the ε_{ran} group to which that observation has been assigned for error analysis—namely, VLT, Keck low contrast (LC), or Keck high contrast (HC). An additional flag highlights two “suspect” absorbers in this catalog—the $z_{\text{abs}} = 1.542$ system toward J00048–415728 and the $z_{\text{abs}} = 2.84$ system toward J194454+770552. These objects were omitted from the Webb et al. team’s most recent work, and so for consistency are omitted from our reanalysis as well. The quasar dataset thus described is available in electronic form from Professor Michael Murphy’s [homepage](#) at the Swinburne University of Technology.

2.1. Explained Errors. An astronomical spectrograph, such as HIRES on Keck or UVES on the VLT, uses a high-resolution diffraction grating to isolate distinct wavelength components of the incoming light, projecting these onto a charge-coupled device (CCD) for digital intensity measurement. The capture of a raw astronomical spectrum thus constitutes a photon counting experiment in which the count in each pixel of the CCD represents a single draw from a Poisson distribution with mean value set by the integrated flux intensity of light within a narrow wavelength interval from the observed source *plus* a contaminating contribution from the background sky and even possibly other non-target emission (e.g. stray light from other bright sources in the field of view); all *modulated* (i.e., non-linearly transformed) by the inherent response characteristic of the telescope and device. In the Webb et al.

team’s analysis multiple such raw spectra for each quasar were sourced from the Keck and VLT archives and *reduced* (i.e., combined and “cleaned”) to single sequences of (high quality) absorption line profiles ready for modeling. The reduction process necessarily involves a number of complex transformations of the raw data (including “stacking”, “flat fielding”, “bias frame subtraction”, “wavelength calibration”, “cosmic ray removal”, and “continuum subtraction”; the essential cleaning and calibration procedures of precision CCD photometry), which ultimately correlate to some extent both nearby and not-so-nearby pixels in the output profile sequence. However, given the almost overwhelming complexity of accounting robustly for each stage of the data reduction pipeline the extent of this correlation is assumed negligible—as is standard practice in the field⁶—with each pixel in the reduced spectrum ultimately treated as independent of all others and assigned an observational standard deviation, σ_i .⁷

As noted earlier the MM method for deriving from the reduced spectrum a $\Delta\alpha/\alpha$ estimate for each intervening absorber involves a complex multi-parameter fit across its signature absorption lines. Due to both natural (collisional) and Doppler (thermal and/or turbulent) broadening, the imprint of each line will be typically spread out over multiple resolution elements and a *profile* function thus required to characterize its shape and recover its centroid (i.e., the precise redshift of that component). The Voigt profile function⁸—characterized by the three controlling parameters: redshift, column density, and velocity width; with the latter constrained jointly (as the kinetic temperature) across all species detected in the thermal broadening case and independent in the turbulent case—is well-established as the natural basis for this procedure. However, the number of such bases required to accurately model a particular system, itself a function of the number of distinct “velocity components” present in the intervening cloud, is of course unknown *a priori*. The Webb et al. team therefore elected to fit each absorber sequentially, starting with a single Voigt profile function for each line and

⁶Substantial research efforts are, however, underway to develop Bayesian techniques for modeling more faithfully the complete observational reduction process; see, for example, the recent thesis on this topic by [Bosch \(2011\)](#).

⁷For further (highly technical) details of this reduction process and the rules applied for “error bar propagation” therein the interested reader may refer, for instance, to the official UVES data reduction cookbook available [online](#).

⁸The Voigt profile function ([Kendall, 1938](#); [Nason, 2006](#)) corresponds in shape to the probability density of a continuous random variable defined as the sum of two independent random variables of standard forms—a Normal and a Cauchy. For a more detailed description of the use of the Voigt profile function in the modeling of astronomical spectra the interested reader may refer to (in particular, Appendix A of) Michael Murphy’s PhD thesis also available [online](#).

adding components until a “statistically acceptable” solution was achieved (King et al., 2012); with the benchmark for acceptability characterized by a $1/\sigma_i^2$ -weighted sum of squared residuals (WSSR) approximately equal to the number of pixels fit minus the number of free parameters (dubbed, “the degrees of freedom”, in their analysis though the fitted model is in fact non-linear). The known limitations of the WSSR as a stopping criterion for model selection (cf. Mallows 1995; Andrae, Schulze-Hartung and Melchior 2010) may well contribute to the unexplained errors of the quasar dataset discussed below.

Of those Voigt profile models (both thermal and turbulent) tested up to this maximum number of components the model with minimum Akaike Information Criterion (corrected for small n) was then selected as the best choice for *refitting* with $\Delta\alpha/\alpha$ (now) a free parameter. Having located (via a downhill gradient-type search) the WSSR-minimizing (likelihood-maximizing) solution to this final model the Webb et al. team estimate the magnitude of their explained error term from the local curvature of the likelihood surface in $\Delta\alpha/\alpha$; adopting the Normal approximation, $\varepsilon_{\text{obs}} \sim \mathcal{N}(0, \sigma_{\text{obs}}^2)$. The validity of this approximation to the shape of the likelihood surface in the stated context has been confirmed via Markov Chain Monte Carlo (MCMC) exploration, as described by King et al. (2010).

2.2. Unexplained Errors. As mentioned in the Introduction this estimate of the uncertainty in $\Delta\alpha/\alpha$ derived during profile fitting and governed by the (photon counting) realization noise of the observed spectrum has been declared insufficient by the Webb et al. team to explain the apparent noise level in the quasar dataset. Numerous *possible* reasons for this have been proposed and discussed at length by Murphy, Webb and Flambaum (2003) and King et al. (2012). These may be divided broadly into two classes: (I) methodological limitations of the fitting procedure rendering inadequate the adopted ε_{obs} distribution, and (II) hitherto unmodeled sources of additional random and/or systematic error. From the first class we note the neglected impact of structural uncertainty in the profile fit with regard to the choice of line broadening mechanism and number of velocity components to include; with perhaps a tendency toward underestimation of the latter owing to the known pitfalls of the WSSR stopping criterion adopted. From the second class we have the chance blending of lines from absorbers at different redshifts, and/or run-to-run errors in the wavelength calibration solution (King et al., 2012). Dedicated investigations of the latter by Griest et al. (2010) and Whitmore, Murphy and Griest (2010) have exposed the presence of intra-order velocity shifts against iodine cell reference spectra imaged on both the

HIRES and UVES spectrographs of sufficient magnitude to perhaps impact on estimates of the fine structure constant; see also the novel asteroid-based calibration study of [Molaro et al. \(2008\)](#).

To account for the inadequacy of ε_{obs} in their analysis the Webb et al. team suppose the existence of a Normally-distributed error term, $\varepsilon_{\text{ran}} \sim \mathcal{N}(0, \sigma_{\text{ran}}^2)$, adding strictly unbiased (zero mean and mode) noise to their $\Delta\alpha/\alpha$ estimates ([Webb et al., 2011](#); [King et al., 2012](#); [Berengut, Kava and Flambaum, 2012](#)). For each of their monopole and dipole hypotheses tested against the quasar dataset the authors constrain the magnitude (*viz.* standard deviation) of this error term, σ_{ran} , through a Least Trimmed Squares (LTS) procedure ([Rousseeuw, 1984](#)) in which both its estimate and the best-fitting set of hypothesis parameters (under a WSSR-based likelihood function including this additional error term) are refined in an iterative manner. For the specific monopole+ $r(z)$ -dipole hypothesis examined herein (Equation 2 below) the authors estimate $\sigma_{\text{ran}} \approx 0.858 \times 10^{-5}$ for their VLT sub-sample (or ε_{ran} group), $\sigma_{\text{ran}} \approx 1.630 \times 10^{-5}$ for the sub-sample of their Keck absorbers with high contrast observations (the Keck HC ε_{ran} group), and $\sigma_{\text{ran}} \approx 0$ for the remainder of their Keck absorbers with low contrast observations (the Keck LC ε_{ran} group).

Hence, under the Webb et al. team's final model for both their explained and unexplained uncertainties the total error in each $\Delta\alpha/\alpha$ estimate is treated as $\varepsilon_{\text{tot}} \sim \mathcal{N}(0, \sigma_{\text{tot}}^2)$ where $\sigma_{\text{tot}}^2 = \sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2$ (with σ_{obs} unique to that absorber and σ_{ran} shared across its ε_{ran} group). After outlining for reference the mathematical form of the monopole+ $r(z)$ -dipole hypothesis below we proceed to examine the σ_{obs} - and σ_{tot} -scaled residuals about its best-fit solution from the [King et al. \(2012\)](#) study as a means of evaluating (qualitatively) this simple error treatment.

2.3. Mathematical Form of the Dipole. The monopole+ $r(z)$ -dipole hypothesis considered by [King et al. \(2012\)](#) and [Berengut, Kava and Flambaum \(2012\)](#) takes the functional form:

$$(2) \quad \Delta\alpha/\alpha_{\text{mod}|\mathbf{x}_i, \boldsymbol{\theta}_m} = m + B \times r(z_i) \times \cos(\phi)[\text{ra}_i, \text{dec}_i, \text{ra}_d, \text{dec}_d]$$

with $\mathbf{x}_i = \{\text{ra}_i, \text{dec}_i, r(z_i)\}$ the vector of explanatory variables for the i th absorber, $\boldsymbol{\theta}_m = \{m, B, \text{ra}_d, \text{dec}_d\}$ a vector of input model parameters, and $\cos(\phi)[\cdot]$ a function returning the cosine of angular separation between the observational sightline and dipole vector. Given right ascension in hours and declination in degrees the latter may be computed as:

$$(3) \quad \cos(\phi) = \sin(\text{dec}_i) \sin(\text{dec}_d) + \cos(\text{dec}_i) \cos(\text{dec}_d) \cos(15 \times [\text{ra}_i - \text{ra}_d]).$$

The lookback distance (cf. [Hogg 1999](#)) for each absorber, $r(z_i)$, under the standard cosmological model (here we adopt $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$) is given (in units of GLyr; Giga-Light-years) by:

$$(4) \quad r(z_i) = 13.98 \times \int_0^{z_i} \left[(1+z') \sqrt{0.3 \times (1+z')^3 + 0.7} \right]^{-1} dz'.$$

For reference, we note that the monopole-only null hypothesis corresponds simply to the case of Equation 2 with $B = 0$ —that is, $\Delta\alpha/\alpha_{\text{mod}}|_{\theta_m=\{m\}} = m$ (a non-zero constant); with $\Delta\alpha/\alpha_{\text{mod}}|_{\theta_m=\{m=0\}} = 0$ its strict null counterpart.

2.4. Possible Non-Normality of Residuals. In the top row of Figure 3 we present histograms of the residuals about the best-fit (i.e., $1/\sigma_{\text{tot}}^2$ WSSR-minimizing) solution to the monopole+ $r(z)$ -dipole hypothesis reported by [King et al. \(2012\)](#), scaled by the magnitude of each absorber’s raw observational error term, σ_{obs} , and subdivided by telescope (or, more precisely, ϵ_{ran} group) and redshift. To limit any sensitivity to the precise functional form for the dipole hypothesis assumed herein we include only those systems lying within $\pm 10^\circ$ of its equator where the contribution of the dipole component is smaller than or equal to the magnitude of the fitted monopole component ($|m| = 0.187 \times 10^{-5}$). If only the explained uncertainties (cf. Section 2.1) represented by $\epsilon_{\text{obs}} \sim \mathcal{N}(0, \sigma_{\text{obs}}^2)$ were in operation here (and the monopole+ $r(z)$ -dipole hypothesis both true and well fit), then the distributions of these scaled residuals should, of course, themselves be standard Normal. However, this seems unlikely to be the case (as acknowledged by the Webb et al. team’s invocation of the unexplained error term) for either the VLT residuals—which exhibit an unusual bimodality—or the Keck residuals—which exhibit a strong negative skewness. These departures from the standard Normal are further emphasized by the Q-Q plots shown in the top-right inset of each panel.

As described in Section 2.2, under the Webb et al. team’s model for the unexplained uncertainties in the quasar dataset the *total* error on each observation remains zero mean Normal, $\epsilon_{\text{tot}} \sim \mathcal{N}(0, \sigma_{\text{tot}}^2)$, but takes a broader standard deviation (at least in the case of the VLT and Keck HC ϵ_{ran} groups) of $\sigma_{\text{tot}} = (\sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2)^{1/2}$. Rescaling the VLT and Keck HC residuals accordingly we recover those histograms shown in the bottom row of Figure 3. While the scaled residuals now bear a greater similarity to the standard Normal, from a critical perspective the rough agreement seen here could hardly be considered reassuring, especially given the lack of any strong justification for the supposed form of ϵ_{tot} . By including here a number of plausible alternatives to the Webb et al. team’s nominal error model in our reanalysis of




FIG 3. *Motivation for our consideration of non-Normal error terms. Each panel contains a histogram of the residuals, $[\Delta\alpha/\alpha - \Delta\alpha/\alpha_{\text{mod}}]$, with respect to the best-fit parameterization of the monopole+ $r(z)$ -dipole hypothesis of King et al. (2012), scaled by the standard deviation of the raw observational error term only in the top row, σ_{obs} , and that of the total error term adopted by Webb et al., $(\sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2)^{1/2}$, in the bottom row. To limit any sensitivity to the precise functional form adopted for the dipole we include only those systems lying within $\pm 10^\circ$ of its equator (VLT sources on the left; Keck on the right), and we further subdivide each sample by redshift at $z = 1.5$. We draw the reader’s attention to departures in a number of these histograms from the standard Normal form that would be expected if the measurement errors were governed entirely by Normal terms of the magnitude assumed in each row; in particular, the bimodal split about zero in the VLT residuals, and the negative skewness in the Keck residuals. We further emphasise these departures from Normality via the Q-Q (residual vs. Normal) plots shown as insets in the top-right corner of each panel; the range here (not marked) is exactly -3 to 3 on each axis.*

the quasar dataset we are able to gauge the influence of the assumed form for the unexplained errors on the apparent dipole significance (cf. Sections 3 to 5).

2.5. Twice-Observed Absorbers. As noted earlier although the Keck and VLT telescopes cover generally well-separated regions of sky—the Hawaii-

based Keck being largely restricted to targets in the northern hemisphere, and the Chile-based VLT restricted to targets in the southern hemisphere—there nevertheless appears in the Webb et al. dataset an eight quasar intersection of sightlines common to both (evident in Figure 2 as the star-shaped symbols arising from the overlap of the up-triangle and down-triangle symbols used respectively for the VLT and Keck datapoints). Among the corresponding population of intervening absorbers thus probed there are eleven twice-observed; that is, detected at matched spectroscopic redshifts in the spectra from both instruments (which feature quite different observational selection functions—owing to the unique resolution and wavelength coverages of each—and hence do not necessarily return *identical* sets of absorbers). The pairwise comparison of Keck and VLT $\Delta\alpha/\alpha$ estimates for these twice-observed systems offers a “first-order” test of the unbiased error hypothesis questioned by Chad Orzel and others skeptical of the Webb et al. team’s conclusions.

The eleven scaled differences, $[\Delta\alpha/\alpha_{\text{Keck}} - \Delta\alpha/\alpha_{\text{VLT}}]/(\sigma_{\text{tot,Keck}}^2 + \sigma_{\text{tot,VLT}}^2)^{1/2}$, so recovered form the ordered list: $\{-1.64, -1.40, -0.97, -0.75, -0.56, -0.52, -0.34, -0.09, 0.17, 0.27, 0.30\}$ (rounded to the 2nd decimal place). The predominance of negative differences observed here (eight of the eleven) hints at the presence of systematic biases in the $\Delta\alpha/\alpha$ values recovered from one or both of these telescopes; and indeed a Wilcoxon signed-rank test (Wilcoxon, 1945) allows rejection of the zero mean difference hypothesis at the 5% significance level ($p = 0.017$). Though admittedly not of overwhelming power due to the small sample size available this result nevertheless serves as further motivation (in addition to the alignment of the alleged dipole’s equator along the coverage overlap of the Keck and VLT telescopes) for the biased error models we consider in our Bayesian reanalysis of the quasar dataset below.

3. Generative Models & Posteriors. Having established the importance of a Bayesian reanalysis of the quasar dataset, and for exploring in this context both unbiased and biased error terms, in the Introduction and Section 2, respectively, we now proceed to outline formally the generative models and priors required for this endeavor.

3.1. Error Models. In the interests of thoroughness we explore *four* alternative forms for the total error term operating in each ϵ_{ran} group; each candidate form allowing a quite different distribution for the unexplained uncertainty component. For reference we denote these, **A**, **B**, **C**, and **D**,

respectively, and define as follows:

- (A) $\varepsilon_{\text{tot}}|\beta_{\text{ran}}, \sigma_{\text{ran}} \sim \mathcal{N}(\beta_{\text{ran}}, \sigma_{\text{obs}}^2 + \sigma_{\text{ran}}^2)$,
- (B) $\varepsilon_{\text{tot}}|\beta_{\text{ran}}, \sigma_{\text{ran}} \sim \mathcal{V}(\beta_{\text{ran}}, \sigma_{\text{obs}}, \sigma_{\text{ran}})$ [$\mathcal{V}$ denotes the Voigt distribution],
- (C) $\varepsilon_{\text{tot}}|\xi_{\text{ran}}, v_{\text{ran}}, w_{\text{ran}} \sim \mathcal{N}_{\text{skew}}(\xi_{\text{ran}}, v_{\text{ran}}w_{\text{ran}}/\sqrt{\sigma_{\text{obs}}^2 + w_{\text{ran}}^2}, \sigma_{\text{obs}}^2 + w_{\text{ran}}^2)$,
- (D) $\varepsilon_{\text{tot}}|\eta_{\text{ran}}, \gamma_{\text{ran}} \sim \begin{cases} \mathcal{N}(-\gamma_{\text{ran}}\sigma_{\text{obs}}, \sigma_{\text{obs}}^2) & \text{with probability } \eta_{\text{ran}}, \\ \mathcal{N}(\gamma_{\text{ran}}\sigma_{\text{obs}}, \sigma_{\text{obs}}^2) & \text{otherwise.} \end{cases}$

Error form **A** corresponds, of course, to the scenario of a Normally-distributed source of measurement noise, $\varepsilon_{\text{ran}} \sim \mathcal{N}(\beta_{\text{ran}}, \sigma_{\text{ran}}^2)$, operating in addition to, and independently of, the explained error, $\varepsilon_{\text{obs}} \sim \mathcal{N}(0, \sigma_{\text{obs}}^2)$. At $\beta_{\text{ran}} = 0$ this equates to the strictly unbiased (i.e., zero mean and mode) error term adopted by [King et al. \(2012\)](#); and we therefore consider this limiting case the *nominal* error form (**A**₀) for our analysis, treating the more general $\beta_{\text{ran}} \neq 0$ case as the default *skeptical* form.

Error form **B**, on the other hand, corresponds to the (undesirable) scenario of a *heavy-tailed* source of Cauchy-distributed measurement noise with location, β_{ran} , and scale, σ_{ran} , i.e., $\varepsilon_{\text{ran}} \sim \mathcal{C}(\beta_{\text{ran}}, \sigma_{\text{ran}})$ where $f_{\mathcal{C}|\beta_{\text{ran}}, \sigma_{\text{ran}}}(x) = 1/\pi \times \sigma_{\text{ran}}/(\sigma_{\text{ran}}^2 + (x - \beta_{\text{ran}})^2)$, operating in addition to, and independently of, the explained error. As noted in our earlier discussion of absorption line modeling (Section 2.1), the Voigt profile function (cf. [Kendall 1938](#); [Nason 2006](#)), $\mathcal{V}$, provides an analytical expression for the density of such a distribution arising from the convolution of a Cauchy and a Normal. Namely,

$$f_{\mathcal{V}|\beta_{\text{ran}}, \sigma_{\text{obs}}, \sigma_{\text{ran}}}(x) = \frac{\text{Re}[\omega(z)]}{\sigma_{\text{obs}}\sqrt{2\pi}}, \quad z = \frac{(x - \beta_{\text{ran}}) + i\sigma_{\text{ran}}}{\sigma_{\text{obs}}\sqrt{2}}, \quad -\infty < x < \infty,$$

where $\omega(\cdot)$ in the above denotes the complex error (or “Faddeeva”) function. By allowing, through error form **B**, for the possibility of heavy-tailed behaviour in the unexplained error term we help to ensure the robustness of our conclusions under the Normal error form (**A**) against the possible presence of extreme outliers in the quasar dataset. Though the mean remains *undefined* for the Cauchy, and thus the Voigt as well, we note that $\beta_{\text{ran}} = 0$ nevertheless gives a zero mode case (**B**₀), which we refer to as “unbiased” (remembering in our subsequent discussion that this is not strictly so) and which we examine separately to the more general (skeptical) $\beta_{\text{ran}} \neq 0$ case.

Error form **C** supposes instead a source of *skew Normal* noise, $\varepsilon_{\text{ran}} \sim \mathcal{N}_{\text{skew}}(\varepsilon_{\text{ran}}, v_{\text{ran}}, w_{\text{ran}}^2)$, operating in addition to, and independently of, the explained error. A tendency toward bias in the wavelength calibration solution for a given ε_{ran} group may be one explanation for such an *asymmetric*

distribution in the uncertainties of the quasar dataset. We note here the definition of the skew Normal with normalized shape parameter, v , scale, w , and location parameter, ξ , in terms of the standard Normal density, $\phi(\cdot)$, and distribution function, $\Phi(\cdot)$, for reference:

$$(5) \quad f_{\mathcal{N}_{\text{skew}}|\xi,v,w^2}(x) = \frac{2}{w}\phi\left(\frac{x-\xi}{w}\right)\Phi\left(\frac{v(x-\xi)}{w\sqrt{1-v^2}}\right), \quad -\infty < x < \infty,$$

with $-1 < v < 1$ and $w > 0$. We note also that the sum of a (zero mean) Normal, $\mathcal{N}(0, \sigma_{\text{ran}}^2)$, and a skew Normal, $\mathcal{N}_{\text{skew}}(\xi_{\text{ran}}, v_{\text{ran}}, w_{\text{ran}}^2)$, is again skew Normal, with parameters as specified in our definition of error form **C** above. This may be verified, of course, by reference to the characteristic function of the skew Normal distribution; according to [Pewsey \(2000\)](#), $\Psi_{\mathcal{N}_{\text{skew}}(\xi,v,w^2)}(t) = \exp(i\xi t - w^2 t^2/2)\{1 + i\tau(vwt)\}$ where $\tau(x) = \int_0^x \sqrt{2/\pi} \exp(u^2/2) du$ for $x > 0$ and $\tau(-x) = -\tau(x)$. We consider once again a so-called “unbiased” (here zero mode, but not mean) case (**C**₀), constructed by solving numerically for ξ_{mode} for each σ_{obs} and $\{v_{\text{ran}}, w_{\text{ran}}\}$ parameter pairing, in addition to the skeptical case of free ξ_{ran} .

Last of all, error form **D** corresponds to a scenario of systematic misestimation in which $\Delta\alpha/\alpha$ is alternately *under*-shot or *over*-shot by $\gamma_{\text{ran}} \times \sigma_{\text{obs}}$, with probability η_{ran} of the former (and $1 - \eta_{\text{ran}}$ of the latter). Such a bimodal error distribution might well arise as the result of routine misidentification of the dominant line broadening mechanism (turbulent or thermal) or the number of velocity components present when modeling the observed absorption profile (cf. Section 2.1); and was suggested by the distribution of residuals around the equator of the proposed dipole for the VLT sample (examined in Section 2.4). In the interests of thoroughness, though perhaps not terminological consistency, we also consider an “unbiased” (here zero mean, but not mode) case of this error form (**D**₀) with fixed $\eta = 0.5$.

In Figure 4 we illustrate the nature and diversity of all four candidate forms (**A**, **B**, **C**, and **D**) for the total error term, ϵ_{tot} , under a variety of arbitrary choices for their governing parameters. As in earlier studies by the Webb et al. team, each of the three ϵ_{ran} groups of the quasar dataset—namely, the VLT sample, the low contrast (LC) Keck sample, and the high contrast (HC) Keck sample—is treated *separately* here for error analysis and assigned one of the above. Throughout we employ the notation, **AAA**, **B₀B₀B₀**, **CAD** and so on, to indicate which error form has been assigned to each—Keck LC, Keck HC, and VLT, in that order. We refer specifically to the combination, **AAA**, as our default skeptical model and to the combination, **A₀A₀A₀**, as the nominal error model. With each candidate form admitting two or three governing parameters we must therefore specify a




FIG 4. *Illustration of the four candidate forms (A, B, C, and D) proposed here for modeling the total error term, ϵ_{tot} , of the quasar dataset. We specify a standard deviation, $\sigma_{\text{obs}} = 1 \times 10^{-5}$, for the explained uncertainty component, $\epsilon_{\text{obs}} \sim \mathcal{N}(0, \sigma_{\text{obs}}^2)$, in these plots, and we indicate the diversity of densities thereby admitted under each corresponding total error term by tracing the output for a variety of arbitrary inputs to its governing parameters (detailed in the top-right legend of each panel). For clarity, the “unbiased” cases shown of these forms (A_0 , B_0 , C_0 , and D_0) are clearly distinguished (via a reduced line thickness) from their general (skeptical) counterparts.*

total of between six and nine priors for a complete *error model*.

3.2. Error Model Priors. In constructing priors on the governing parameters of our error models we are guided by the logical observation that the magnitude of the unexplained error term must be of roughly the same order as the explained error term in this dataset ($\overline{\sigma_{\text{obs}}} \approx 2 \times 10^{-5}$); if it were very much smaller it would presumably have gone unnoticed, while if it were very much larger the Webb et al. team would have been unlikely to have proceeded to hypothesis testing without a better understanding of its origin. The same follows from inspection of the σ_{ran} -scaled residuals along the equatorial strip shown in Figure 3. In addition, when using biased forms for

ε_{tot} (cf. Section 3.1 above) we specify *a priori* that any bias in the Keck (LC and HC) error terms must be *negative* and any bias in the VLT error term *positive* as per the skeptical interpretation of the quasar dataset. [Note that we also restrict the prior range of the dipole vector to southern hemisphere pointings in the biased error case in accordance with the rival hypothesis—that there remains a similarly oriented dipole field driving the apparent North–South trend in $\Delta\alpha/\alpha$; cf. Section 3.3 below.] Thus, for the parameters, β_{ran} and σ_{ran} , in our default skeptical error form (**A**), which govern the mean/mode and standard deviation, respectively, of the unexplained term in this case, we suppose $\pi(\beta_{\text{ran}}) \sim \mathcal{N}_{\text{half}}(0, [0.5 \times 10^{-5}]^2)$ and $\pi(\sigma_{\text{ran}}) \sim \text{U}(0, 5 \times 10^{-5})$ —with the half Normal prior on β_{ran} positive for the VLT ε_{ran} group and negative for the Keck LC and HC ε_{ran} groups.

To promote fairness in the marginal likelihood-based comparison of each model–hypothesis pairing we have attempted to make our priors on the parameters of the remaining three error forms as similar as possible to those of form **A**. Thus, for error form **B** we suppose $\pi(\beta_{\text{ran}}) \sim \mathcal{N}_{\text{half}}(0, [0.5 \times 10^{-5}]^2)$ for the mode of the underlying Cauchy (with the same choice of sign for each ε_{ran} group), but allow a (slightly) larger range of $\pi(\sigma_{\text{ran}}) \sim \text{U}(0, \sqrt{2 \log(2)} \times 5 \times 10^{-5})$ on its scale parameter to match our effective prior on the full width at half maximum (FWHM) of this error form to that of **A** (the standard deviation being undefined for the Cauchy, and thus not available for comparison). For error form **C** we begin by setting our prior on the *unnormalized* skew parameter, $v_{\text{ran}}/\sqrt{1 - v_{\text{ran}}^2}$, of this distribution to a generous, $\mathcal{N}(0, [10]^2)$; which transforms to the following density on the *normalized* skew parameter, v_{ran} , with $a = 10$:

$$f_{\pi(v_{\text{ran}}, a)}(x) = \frac{1}{\sqrt{2\pi a^2}} \exp\left[\frac{-x^2}{2a^2(1 - x^2)}\right] \frac{1}{(1 - x^2)^{3/2}}, \quad -1 < x < 1.$$

We then match our prior on the standard deviation of the unexplained error term in **C** to that of **A** by setting $\pi(w_{\text{ran}}|v_{\text{ran}}) \sim \text{U}(0, (1 - \frac{2}{\pi}v_{\text{ran}}^2)^{-1/2} \times 5 \times 10^{-5})$; and likewise to match our priors on the corresponding means we choose, $\pi(\xi_{\text{ran}}|v_{\text{ran}}, w_{\text{ran}}) \sim \mathcal{N}_{\text{half}}(-v_{\text{ran}}w_{\text{ran}}\sqrt{\frac{2}{\pi}}, [0.5 \times 10^{-5}]^2)$ signed as above. Finally, for error form **D**, for which no precise matching of this nature is possible, we simply adopt $\pi(\gamma_{\text{ran}}) \sim \text{U}(0, 2.5)$ and $\pi(\eta_{\text{ran}}) \sim \text{Beta}_{\text{half}}(15, 15)$ (i.e., “folded” about 0.5 [$f_{0 < x < 0.5}^* = f_{0.5 < x < 1}^* = 2f_{0 < x < 1}$] with $\eta_{\text{ran}} > 0.5$ assumed for the Keck ε_{ran} groups and $\eta_{\text{ran}} < 0.5$ for the VLT ε_{ran} group), favoring a near-symmetric ε_{ran} over a markedly asymmetric one. For the typical absorber with explained error, $\sigma_{\text{obs}} \approx 2 \times 10^{-5}$, this gives a prior expectation on the bias of the unexplained error term in each ε_{ran} group comparable to that of our form **A** benchmark.

3.3. Hypothesis Priors. We apply similar principles with regard to constructing priors on the input parameters, $\theta = \{m, B, \text{ra}_d, \text{dec}_d\}$, of the Webb et al. team’s proposed monopole+ $r(z_i)$ -dipole hypothesis for the apparent cosmic variation of $\Delta\alpha/\alpha$ (see Equation 2 in Section 2.3). For the strength of the monopole term, m , we adopt a zero mean Normal prior with standard deviation, 0.5×10^{-5} , and for the strength of the dipole term, B , an exponential prior with rate, $1/[0.5 \times 10^{-5}]$. That is, $\pi(m) \sim \mathcal{N}(0, [0.5 \times 10^{-5}]^2)$ and $\pi(B) \sim \text{Exp}(1/[0.5 \times 10^{-5}])$. Supposing *a priori* for all “unbiased” error models that the dipole vector might point anywhere on the celestial sphere with equal probability we assign the appropriate uniform priors to both ra_d and the sine of dec_d . Thus, $\pi(\text{ra}_d) \sim \text{U}(0, 24)$ and $\pi(\sin(\text{dec}_d)) \sim \text{U}(-1, 1)$, i.e., $\pi(\text{dec}_d) \sim \cos(\text{dec}_d)/2$. While for the general (skeptical) case of each model we restrict the dipole declination to the southern hemisphere (i.e., $\text{dec}_d < 0$ with $\pi(\text{dec}_d) \sim \cos \text{dec}_d$) as noted earlier.

For consistency with the above we also suppose $\pi(m) \sim \mathcal{N}(0, [0.5 \times 10^{-5}]^2)$ for the strength of the monopole in the monopole-only version of the null hypothesis; recall though, our strict null hypothesis is that $\Delta\alpha/\alpha$ remains everywhere zero.

3.4. Likelihood Function. With any stochastic variation in the $\Delta\alpha/\alpha$ estimates of the quasar dataset assumed to arise solely from the combination, ε_{tot} , of the explained and unexplained error terms described by the error models of Section 3.1—and with the realization of this measurement error for any given absorber assumed independent of all others—the likelihood function for the observed data, $\mathbf{y} = \{\Delta\alpha/\alpha_i : i = 1, \dots, 293\}$ with covariates represented as $\mathbf{x}_i = \{\text{ra}_i, \text{dec}_i, r(z_i)\}$, corresponding to a given $\Delta\alpha/\alpha_{\text{mod}}$ and ε_{tot} pairing, M , may be written as:

$$(6) \quad L(\mathbf{y}|\theta = \{\theta_m, \theta_e\}, M, \mathbf{x}_i) = \prod_{i=1}^{293} f_{\varepsilon_{\text{tot}}|\theta_e, \varepsilon_{\text{ran}, i}}(\Delta\alpha/\alpha_i - \Delta\alpha/\alpha_{\text{mod}}|\mathbf{x}_i, \theta_m).$$

Here $\theta_m = \{m, B, \text{ra}_d, \text{dec}_d\}$ (dipole) [or $\{m\}$ (monopole null), or $\{m = 0\}$ (strict null)] denotes the set of input hypothesis parameters, $f_{\varepsilon_{\text{tot}}|\theta_e, \varepsilon_{\text{ran}, i}}$ the ε_{tot} probability density for the particular ε_{ran} group to which the i th absorber belongs, and θ_e the vector of between six and nine parameters required for a complete error model.

3.5. Posteriors. For each combination of error model tested here (see Section 4 below for a complete enumeration) and hypothesis (strict null, monopole null, and dipole) we have constructed a Markov Chain Monte

Carlo sample of 100,000 draws from the posterior using a symmetric perturbation, $\boldsymbol{\theta}_{\text{prop}} \sim \mathcal{N}(\boldsymbol{\theta}_i, \boldsymbol{\Sigma}_{\text{prop}})$, for the proposal. The diagonal variance matrix, $\boldsymbol{\Sigma}_{\text{prop}}$, of which was initialized to our prior variance on each parameter and progressively rescaled toward a target acceptance rate of 0.4 during a 5,000 draw burn-in phase. For each final output sample the corresponding autocorrelation function has been computed to verify convergence and an effective sample size of $>10,000$. Although the *marginal likelihoods* of each model–hypothesis pairing—which we derive only indirectly from these posteriors (see Section 4 below)—are in fact the sole quantities of interest for our Bayesian assessment of the claimed spatial variation in the quasar dataset we nevertheless review briefly here the nature of our posterior inferences for key error model and hypothesis parameters as an aid to understanding the ensuing results.

In Figures 5 and 6 we illustrate the marginal posterior probability densities of those parameters (or combinations thereof) governing the width (*viz.* standard deviation, or half width half maximum as appropriate in the case of model **B**) resulting from each error form ($\mathbf{A}/\mathbf{A}_0$ to $\mathbf{D}/\mathbf{D}_0$, as described in Section 3.1) applied homogeneously across all ε_{ran} groups under both the strict null and dipole hypotheses. (Our posteriors under the monopole-only null hypothesis, and for the two non-homogeneous error model combinations we test, **CAA** and **CAD** [cf. Section 4 below], being roughly intermediate to these representative examples.) Although the inferred width of the unexplained error term for each ε_{ran} group varies somewhat depending upon the error model adopted, the relative ordering of the aforesaid—Keck LC, VLT, Keck HC (from smallest to largest)—does not. For the “unbiased” case of each model in particular (Figure 6: $\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$ to $\mathbf{D}_0\mathbf{D}_0\mathbf{D}_0$) we note that the error widths recovered under the null hypothesis are generally somewhat greater than those recovered under the (more flexible) dipole hypothesis—that is, the proposed error forms must necessarily “stretch” to encompass the broader spread of residuals in the former. Interestingly, the Bayesian analysis we present here for the nominal error model ($\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$) favors a non-zero σ_{ran} of roughly 0.6×10^{-5} (posterior mean) for the Keck LC ε_{ran} group under the strict null hypothesis, in contrast to the $\sigma_{\text{ran}} \approx 0$ Least Trimmed Squares (LTS) estimate reported by King et al. (2012) in their Table 2. This discrepancy presumably arises from the deliberate focus on interquartile width at the expense of (or, purportedly in robustness against) any marked outliers in the latter approach. Our estimates for the other ε_{ran} groups under both the null and dipole hypotheses are nevertheless in fair agreement with their LTS counterparts.

In Figure 7 we present the marginal posterior probability densities of the




FIG 5. Posterior probability densities for the key parameters (or combinations thereof) governing the underlying error distribution width (*viz.* standard deviation, or half width half maximum as appropriate in the **BBB** case) of the ϵ_{ran} component in each of our general (skeptical) error models: **AAA** to **DDD** (homogeneous). In the interests of illustrative clarity only those posteriors corresponding to the strict null (*i.e.*, $\Delta\alpha/\alpha$ everywhere zero) and dipole hypotheses are compared here (our results for the monopole-only null hypothesis being roughly intermediate to these two). Interestingly, although the inferred width of the unexplained error term for each ϵ_{ran} group varies somewhat depending upon the error model adopted, the relative ordering of the aforesaid—Keck LC, VLT, Keck HC (from smallest to largest)—does not.

parameters (or combinations thereof) governing the bias (*viz.* offset of the mode, or in case **D** the mean, from zero) in each of our general (skeptical) error models (applied homogeneously across all ϵ_{ran} groups) under both the strict null and dipole hypotheses. As explained in Section 3.2 the posteriors for the Keck (LC and HC) and VLT ϵ_{ran} groups are ascribed priors restricting the sign of bias to negative in the former and positive in the latter in line with the skeptical interpretation of the quasar dataset; hence the solid black divides marked at zero on this Figure. The degree of bias favored by these posteriors for each ϵ_{ran} group varies little between error models but




FIG 6. Posterior probability densities for the key parameters (or combinations thereof) governing the underlying error distribution width (viz. standard deviation, or half width half maximum as appropriate in the $B_0B_0B_0$ case) of the ϵ_{ran} component in each of our “unbiased” error models: $A_0A_0A_0$ to $D_0D_0D_0$ (homogeneous). Again, although the inferred width of the unexplained error term for each ϵ_{ran} group varies somewhat depending upon the error model—and hypothesis—adopted, the relative ordering of the aforesaid—Keck LC, VLT, Keck HC (from smallest to largest)—does not. As expected the error widths recovered under the null hypothesis here are generally somewhat greater than those recovered under the (more flexible) dipole hypothesis.

greatly between hypotheses—as expected, larger biases are generally preferred under the strict null hypothesis than under the dipole hypothesis, owing to the inherent (though of course only partial) degeneracy between the North–South dipole and opposing bias scenarios.

Finally, for reference we plot in Figure 8 the joint posterior densities of the key parameter pairings under the monopole+ $r(z)$ -dipole hypothesis—viz. monopole strength–dipole strength (m – B) and right ascension–declination of the dipole vector (ra_d – dec_d)—for error models AAA , $A_0A_0A_0$, CCC , and $C_0C_0C_0$. While the posteriors corresponding to each error model in




FIG 7. Posterior probability densities for the key parameters (or combinations thereof) governing the error distribution bias (viz. offset of the mode, or in case **D** the mean, from zero) in each of the biased error models: **AAA** to **DDD** (homogeneous). Once again in the interests of illustrative clarity only those posteriors corresponding to our strict null hypothesis and dipole model are compared here. As explained in Section 3.2 the Keck (LC and HC) and VLT ϵ_{ran} groups are ascribed priors restricting the sign of their bias to negative in the former and positive in the latter in line with the skeptical interpretation of the quasar dataset; hence the solid black divides marked here at zero. The degree of bias favored by the results for each ϵ_{ran} group varies little between error models but greatly between hypotheses—as expected, much larger biases are inferred under the strict null hypothesis than under the dipole hypothesis, owing to the inherent (though of course only partial) degeneracy between the North–South dipole and opposing bias scenarios.

the “unbiased” case are both highly concentrated around the maximum likelihood solutions identified by the Webb et al. team under their fitting of the nominal error model ($\mathbf{A}_0 \mathbf{A}_0 \mathbf{A}_0$)—namely, $\{\text{ra}_d \approx 17.5 \text{ hours}, \text{dec}_d \approx -60 \text{ degrees}\}$ and $\{m \approx -0.2 \times 10^{-5}, B \approx 10^{-6}\}$ —in the biased (skeptical) case the corresponding posteriors are evidently more diffuse and favor markedly a smaller dipole strength.

To discriminate between this plethora of plausible generative models with




FIG 8. *Posterior probability densities for the four key parameters, $\theta_m = \{m, B, ra_d, dec_d\}$, governing the Webb et al. team’s proposed monopole+ $r(z)$ -dipole hypothesis for the apparent spatial variation of $\Delta\alpha/\alpha$ in the fine structure dataset under the following alternative, example forms of the unexplained error term: **AAA** and **A₀A₀A₀** (top row); and **CCC** and **C₀C₀C₀** (bottom row). We illustrate each posterior via a series of smoothed contours (surrounding the posterior mode and) enclosing increasing fractions of the (marginal) posterior mass from 0.05 to 0.95 in both the m – B plane (giving the strengths of the monopole and dipole components, respectively) and the ra_d – dec_d plane (giving the dipole direction on the celestial sphere). While the posteriors under each error model in the “unbiased” case are both highly concentrated around the maximum likelihood solutions identified by the Webb et al. team under their fitting of the nominal error model (**A₀A₀A₀**)—namely, $\{ra_d \approx 17.5 \text{ hours}, dec_d \approx -60 \text{ degrees}\}$ and $\{m \approx -0.2 \times 10^{-5}, B \approx 10^{-6}\}$ —in the biased (skeptical) case the corresponding posteriors are evidently more diffuse and favor markedly a smaller dipole strength.*

quasi-degenerate parameters—and to thereby identify the strongest candidate(s) for explaining the apparent spatial variation of the quasar dataset—we now proceed to the Bayesian comparison of marginal likelihoods (cf. [Carlin and Chib 1995](#); [Kass and Raftery 1995](#)).

4. Marginal Likelihoods. For each combination of proposed error model and hypothesis we compute the corresponding marginal likelihood via both *nested importance sampling* (Chopin and Robert, 2010) and the method of *power posteriors* (Friel and Pettitt, 2008)—the former allowing the sensitivity of the resulting Bayes factors to our chosen priors to be rapidly assessed, and the latter providing an important verification of our evidence computations.

4.1. *Nested Importance Sampling.* The marginal likelihood, Z , of the observed data, $\mathbf{y}$, under a given model, M , is (most generally) defined as the integral of the corresponding likelihood function, $L(\mathbf{y}|\boldsymbol{\theta}, M)$, over the prior density, $\pi(\boldsymbol{\theta}|M)d\boldsymbol{\theta}$; thus,

$$(7) \quad Z = \int_{\Omega(\boldsymbol{\theta})} L(\mathbf{y}|\boldsymbol{\theta}, M)\pi(\boldsymbol{\theta}|M)d\boldsymbol{\theta}.$$

With $\boldsymbol{\theta}$ typically of high dimension, and the bulk of the likelihood concentrated over a small volume of the prior domain, in practical applications one cannot hope for an accurate evaluation of Z via direct numerical integration (*viz.* quadrature); so computational MC methods are most commonly relied upon for this task (see, e.g., Friel and Wyse 2012 for a recent overview). Of these MC methods, nested sampling (Skilling, 2006), is perhaps the most popular among the physics and cosmology community at present (e.g. Mukherjee, Parkinson and Liddle 2006; Feroz, Hobson and Bridges 2009), owing to its surprising efficiency for cases in which the prior can be analytically transformed to an n -dimensional Uniform density.⁹

Nested sampling aims to estimate the marginal likelihood via a transformation of the variable of integration in Equation 7 to the prior mass cumulant differential, dX , defined

$$(8) \quad X(\lambda) = \int_{L(\mathbf{y}|\boldsymbol{\theta}, M) > \lambda} \pi(\boldsymbol{\theta}|M)d\boldsymbol{\theta}.$$

As such, $X(\lambda)$ represents the prior probability of the system parameters lying within that restricted domain, $\{\boldsymbol{\theta} : L(\mathbf{y}|\boldsymbol{\theta}, M) > \lambda\}$, for which the

⁹For instance, in cosmological applications the prior is typically assumed independently Uniform on each separate model parameter or the logarithm thereof. This allows the very fast (with respect to the MCMC evolution alternative) ellipse-based strategy of Mukherjee, Parkinson and Liddle (2006) to be applied at the constrained-likelihood sampling phase crucial to this algorithm (see below). No such transformation is possible for our priors on the controlling parameters of error model $\mathbf{C}$ in this study, however, as we have analytical forms available only *conditionally* for $w_{\text{ran}}|v_{\text{ran}}$ and $\xi_{\text{ran}}|v_{\text{ran}}, w_{\text{ran}}$; and likewise, for instance, in the case of the Normal-Gamma prior considered by Friel and Wyse (2012) in their review.

corresponding likelihood exceeds the input, λ ; and as a probability its range is simply zero to one. Supposing the existence of an inverse-defined function for the likelihood, $L(X(\lambda)) \equiv \lambda$, (i.e., supposing $L(\mathbf{y}|\boldsymbol{\theta}, M)$ continuous and $\Omega(\boldsymbol{\theta})$ connected, cf. [Chopin and Robert 2010](#)) the nested sampling formulation of Equation 7 under dX becomes simply

$$(9) \quad Z = \int_0^1 L(X) dX.$$

As [Skilling \(2006\)](#) demonstrates, through the maintainance of a reference particle population sampled from the prior under a non-decreasing likelihood constraint one may generate a sequence of cumulant–likelihood pairings, $\{X_i, L_i\}$, with each L_i known precisely and the corresponding X_i sequence known at least in distribution (decreasing near-geometrically toward zero). Given a sample from, or asymptotic sampling approximation to, this baseline X_i sequence the Z in Equation 9 may be conveniently approximated via a simple Riemann sum or trapezoidal rule (cf. the procedure given in Section 6 of [Skilling 2006](#)); with a dominant stochastic error—easily estimable by repeated recomputation of Z under resampled X_i —for this process of order $O(N^{-1/2})$ governed by the size, N , of the reference population ([Chopin and Robert, 2010](#)).

The computational efficiency of the nested sampling algorithm in practical applications will most often be limited by one’s ability to sample from the prior under a strict likelihood constraint. Unless the prior can be transformed to an n -dimensional Uniform density as noted above it will be necessary to run an MCMC chain in order to explore (and thereby sample “unbiasedly” from) this constrained volume, starting from the position of an existing reference particle and finishing after sufficient moves to obtain approximate stationarity ([Skilling, 2006](#); [Friel and Wyse, 2012](#)); resulting in an overhead of perhaps ~ 20 likelihood evaluations per $\{X_i, L_i\}$ pairing. As a means to take advantage of the accuracy provided by the geometric exploration of the cumulant baseline in nested sampling, while side-stepping the implementational difficulties of constrained likelihood exploration, [Chopin and Robert \(2010\)](#) have proposed *nested importance sampling*. In this alternative approach the true posterior is approximated via an instrumental prior, $\tilde{\pi}(\boldsymbol{\theta})$, and instrumental likelihood, $\tilde{L}(\boldsymbol{\theta})$, with forms chosen to facilitate constrained likelihood sampling; and a weighted sum of the true likelihood function and true prior at each drawn $\boldsymbol{\theta}_i$ is then used to approximate Z .

As in Section 5 of [Chopin and Robert \(2010\)](#) we adopt here a multivariate Normal form for $\tilde{\pi}(\boldsymbol{\theta})$ with mean vector, $\tilde{\boldsymbol{\mu}}$, and covariance matrix, $\tilde{\boldsymbol{\Sigma}}$, set to their sampling estimates from the MCMC output of each model–hypothesis combination tested. Supposing an arbitrary, decreasing function

in quadratic θ for $\tilde{L}(\theta)$ thus allows the required sequence of θ_i corresponding exactly to the nested sampling X_i under N reference particles to be generated from $\theta_i = \tilde{\mu} + q_i^{1/2} \mathbf{C} \mathbf{v} / \|\mathbf{v}\|_2^{1/2}$, where $\mathbf{C}$ is the Cholesky lower triangle of $\tilde{\Sigma}$, $\mathbf{v} \sim \mathcal{N}(0, \mathbf{I})$, and $q_i = Q_{\chi_d^2}(e^{-i/N})$ [here $Q_{\chi_d^2}$ represents the quantile function of the χ^2 distribution of degree, d]. Both d and the dimension of $\mathbf{I}$ in this expression are, of course, set equal to the dimension of θ . By running this nested importance sampling procedure on each of our error model–hypothesis pairings with $N = 500$ and the end point of summation set to well beyond the point of diminishing returns at $5N$ iterations we find that the desired marginal likelihoods can be recovered to within an accuracy of $\Delta \log Z = \pm 0.1$ [standard error from repeat simulation] in negligible computational time—in particular, much faster than the power posteriors method—with one exception, the case of the dipole hypothesis fit under error model $\mathbf{C}_0 \mathbf{C}_0 \mathbf{C}_0$.

The reason for this exception is that the posterior for this particular case is distinctly multimodal—featuring near equal mass peaks at positive and negative values of the normalized shape parameter for each ε_{ran} group—meaning that our multivariate Normal instrumental prior is no longer a reasonable approximation. In principle it may be possible to extend nested importance sampling to the multimodal case through a strategy akin to multimodal nested sampling (Feroz and Hobson, 2007). However, for the purposes of the present analysis we rely solely upon our power posteriors-based estimate of the marginal likelihood for this specific error model–hypothesis pairing ($\mathbf{C}_0 \mathbf{C}_0 \mathbf{C}_0$ under the dipole)—the power posteriors approach being applicable to a wider class of distributions including such multimodal forms.

Finally, we note that by recording the sequence of θ_i drawn along with each $\{X_i, L_i\}$ pairing and $\tilde{\pi}(\theta_i)$ density under nested importance sampling one may quickly recompute $\log Z$ in response to any “compatible” modification to the prior density—“compatible” meaning here *technically* that the support of the new prior is contained within that of the old, and *practically* (for error control purposes) that this modification does not too drastically reduce the effective sample size of the weighted sum. We specifically exploit this property of nested importance sampling in the present study (see Section 4.3) in order to check the sensitivity of certain key results to our chosen priors, thereby addressing to some extent potential concerns regarding subjectivity in this aspect of Bayesian model selection (cf. Chipman, George and McCulloch 2001).

4.2. Power Posteriors. The power posteriors method for evaluating the marginal likelihood (cf. Friel and Pettitt 2008) takes a thermodynamic inte-

gration/path sampling approach (Gelman and Meng, 1998; Lefebvre, Steele and Vandal, 2010) to this normalization problem inspired by the methodology of theoretical physics. The power posterior at a given “temperature”, t , is defined as the distribution:

$$(10) \quad \pi(\boldsymbol{\theta}|\mathbf{y}, M, t) \propto L(\mathbf{y}|\boldsymbol{\theta}, M)^t \pi(\boldsymbol{\theta}|M),$$

with normalization (in general) unknown except at $t = 0$. As demonstrated by Friel and Pettitt (2008) the direct marginal likelihood integral of Equation 7 may be reformulated as an integral of the mean log likelihood over the family of power posteriors spanning $t = 0$ to 1; that is,

$$(11) \quad \log Z = \int_0^1 \left\{ \int_{\Omega(\boldsymbol{\theta})} [\log L(\mathbf{y}|\boldsymbol{\theta}, M)] \pi(\boldsymbol{\theta}|\mathbf{y}, M, t) d\boldsymbol{\theta} \right\} dt.$$

Thus, via MCMC sampling from each power posterior over a prescribed temperature schedule—representing (for the typical case of $t_0 = 0 < t_1 < \dots < t_{n-1} < t_n = 1$) a thermodynamic integration from the prior to the posterior—one can approximate the $\log Z$ of Equation 11 using (again) a Riemann sum or trapezoidal rule applied to the corresponding sequence of approximate mean log likelihoods. Here we apply a heating schedule of 40 increments uniformly-spaced in $\sqrt{t}$ and draw 10,000 log likelihoods (after a 1,000 step burn in phase) from each power posterior (to estimate $\langle \log L \rangle_{\pi_t}$).

4.3. Results. Marginal likelihoods were computed for ten alternative permutations of our four candidate error forms (Section 3.1) over the three ϵ_{ran} groups of the quasar dataset—in particular, the four general (skeptical) error forms applied homogeneously (**AAA**, **BBB**, **CCC**, **DDD**), their “unbiased” case counterparts applied likewise (**A₀A₀A₀**, **B₀B₀B₀**, **C₀C₀C₀**, **D₀D₀D₀**), plus two non-homogeneous permutations (**CAD** and **CAA**) motivated by the observed distributions of residuals ($\Delta\alpha/\alpha - \Delta\alpha/\alpha_{\text{mod}}$) about the King et al. (2012) dipole fit along the equatorial strip of Figure 3—under each of the three competing hypotheses—namely, strict null, monopole-only null, and monopole+ $r(z)$ -dipole. These results, totalling thirty separate marginal likelihood estimates, are compiled (in $\log Z$ format) in Table 1.

The first point we note here upon consideration of the above is that for the nominal error model (**A₀A₀A₀**) of the Webb et al. team—in which the unexplained source of error in each ϵ_{ran} group is assumed strictly Normal and strictly unbiased (zero mean and mode)—the log marginal likelihood of the dipole hypothesis ($\log Z = -612.2$) exceeds that of both the strict null ($\log Z = -617.5$) and monopole-only null ($\log Z = -617.4$) alternatives by

TABLE 1
Log Marginal Likelihoods for Each Error Model–Hypothesis Pairing Tested Against the Quasar Dataset

	AAA	A ₀ A ₀ A ₀	BBB	B ₀ B ₀ B ₀	CCC
Strict Null	-607.8	-617.5	-619.3	-628.6	-608.7
Monopole	-608.4	-617.4	-620.0	-628.4	-609.7
Dipole	-609.3	-612.2	-620.2	-622.4	-609.9
	C ₀ C ₀ C ₀	DDD	D ₀ D ₀ D ₀	CAD	CAA
Strict Null	-617.1	-623.4	-627.6	-614.8	-607.8
Monopole	-615.1	-624.0	-628.4	-615.4	-607.9
Dipole	-612.4	-616.2	-618.4	-616.2	-608.5

$\Delta \log Z \approx 5.2$. Under the Jeffreys scale for interpretation of *Bayes factors* (B.F. = $\exp[\Delta \log Z]$) such an extreme result (a B.F. > 100) should be considered *decisive* evidence in favor of the former; broadly consistent *at face value* (i.e., neglecting for now both our results for alternative error models and any subjective priors one might hold on the relative merits of these hypotheses) with the Webb et al. team’s bootstrap randomization-based estimate of $\sim 4\sigma$ support for spatial variation in the quasar dataset. [We say “broadly” as of course: (i) one cannot meaningfully compare p -values against Bayes factors in a quantitative sense; and (ii) a log marginal likelihood offset of $\Delta \log Z \approx 5.2$ would in any case correspond only to $\approx 2.5\sigma$ support on the one-sided Normal scale.]

A further inspection of the results compiled in Table 1 reveals that this particular ordering of the three hypotheses by marginal likelihood recovered under the nominal error model ($\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$) is in fact conserved across all three alternative error models tested in their “unbiased” cases—namely, $\mathbf{B}_0\mathbf{B}_0\mathbf{B}_0$ (heavy tailed), $\mathbf{C}_0\mathbf{C}_0\mathbf{C}_0$ (skew), and $\mathbf{D}_0\mathbf{D}_0\mathbf{D}_0$ (bimodal). But, one may also note that the greatest marginal likelihood recovered here for the (*prima facie*) preferred dipole hypothesis under any of these error models remains that of $\log Z = -612.2$ for the nominal error model (unbiased, Normal noise); under the interpretation of the Bayes factor as Ockham’s razor (cf. [Jeffreys and Berger 1992](#); [Jaynes 2003](#)) we must conclude that there exists at present insufficient evidence to favor any of these more complex error models over the nominal in their “unbiased” forms.

Turning now to the results in Table 1 for our general (skeptical) error models, which expressly allow for the possibility of opposing systematic biases between the observations from the Keck and VLT telescopes, we discover for all but one permutation tested ($\mathbf{DDD}$) a reversal of the above rank-

ing of hypotheses; that is, for error models **AAA**, **BBB**, **CCC**, **CAD**, and **CAA** the strict null and monopole-only null hypotheses are in fact preferred in marginal likelihood over the Webb et al. team’s monopole+ $r(z)$ -dipole hypothesis—although we note that the strength of this preference ranges only between “barely worth mentioning” and “substantial” on the Jeffreys scale. Furthermore, for three of these error models (**AAA**, **CCC**, and **CAA**) the resulting marginal likelihoods for all hypotheses are consistently greater than the maximum recovered for the dipole hypothesis under the nominal error model; with a tie for the greatest maximum likelihood of the present study ($\log Z = -607.8$) between **AAA** and **CAA** under the strict null hypothesis. That we recover such support for a skew Normal term in the Keck LC ε_{ran} group is quite intriguing in that it may offer some insight into the origin of the unexplained errors of this dataset—in particular, it suggests a sole contributor, such as bias in the Keck wavelength calibration solution, rather than the accumulation of many disparate sources converging toward a Normal. Nevertheless, we have elected to focus the ensuing discussion on the tied **AAA** model as this was our *a priori* default skeptical proposal and is most similar in form to its (unbiased) nominal rival.

As mentioned earlier, in order to investigate the sensitivity of these results to our choice of priors on both error model and hypothesis parameters we make use here of an important property of the nested importance sampling algorithm—namely, that it allows for rapid recomputation (i.e., requiring no new likelihood evaluations) of the marginal likelihood under (“compatible”) modifications to the prior density. In Figure 9 we explore the subjectivity in this regard of our recovered Bayes factors (B.F.) comparing the monopole+ $r(z)$ -dipole hypothesis against the strict null. Specifically, in the left hand panel we illustrate the effect of modifying our prior “standard deviation” (σ_p ; the quotation marks referencing the fact that the σ_p hyper-parameter described here is not the true standard deviation for this *half Normal* prior) on the bias (β_{ran}), and our prior range (u_q) on the width (σ_{ran}), of the ε_{ran} component in our default skeptical error model (**AAA**); and we do the same for our prior standard deviation (σ_m) on the monopole strength term (m), and our prior rate (u_n) on the dipole strength term (B), of the dipole hypothesis in the right hand panel. In each case it is evident that large modifications of our stated prior beliefs are required to overturn the weight of evidence against spatial variation in the fine structure constant under this biased error model—either a drastic rescaling of the prior density on the magnitude of this bias toward zero (i.e., moving back toward the unbiased alternative), or a significant tightening of our prior on the dipole strength term (bringing it toward its posterior mean, though we have no *a*




FIG 9. *Exploring the sensitivity to our chosen priors of the Bayes factor (B.F.) comparing the monopole+ $r(z)$ -dipole hypothesis against the strict null under the default skeptical error model (AAA). By re-weighting the summation of $\{X_i, L_i\}$ pairings under each new $\pi(\theta_i)$ in the nested importance sampling computation of $\log Z$ for each hypothesis we explore in the lefthand panel the effect of varying our prior “standard deviation” (σ_p) on the bias (β_{ran}), and prior range (u_q) on the width (σ_{ran}), of the unexplained errors (ϵ_{ran}); and we do the same for our prior standard deviation (σ_m) on the monopole strength term (m), and our prior rate (u_n) on the dipole strength term (B), of the dipole hypothesis in the righthand panel. In each case it is evident that large modifications of our stated prior beliefs (indicated by the purple asterix overlaid on each plot) are required to overturn the weight of evidence against spatial variation in $\Delta\alpha/\alpha$ under this biased error model. (Note that the Bayes factors for $u_q > 5 \times 10^{-5}$ shown in the lefthand panel are shaded differently to indicate that the adopted rescaling beyond this point is not technically “compatible” in the sense that the new prior support is not contained within that of the old; but this should not appreciably affect the accuracy of our marginal likelihood evaluations presented here as the likelihood for all $\sigma_{\text{ran}} > 5 \times 10^{-5}$ is itself exceedingly small.)*

priori theoretical motivation for such a small dipole strength¹⁰).

In summary, the marginal likelihood comparisons presented herein point toward the following key result: *that the Webb et al. team’s claimed “ 4σ support” for a dipole signature in the quasar dataset depends crucially upon their assumption that the unexplained error terms in the Keck and VLT subsamples contribute only **unbiased** noise to the estimation of $\Delta\alpha/\alpha$.* We now discuss these results more fully (i.e., in the context of our subjective priors on the plausibility of each error model–hypothesis pairing) with regard to the evidence, or lack thereof, for a spatially-varying fine structure constant in Section 5 below.

¹⁰Indeed, Sean Carroll has argued (in his blog post mentioned in the Introduction) that *if anything* one would expect a far *greater* dipole strength considering the energy level implied for the underlying scalar field.

5. Discussion and Conclusions. Although (as mentioned in the Introduction) there exist various exotic physical theories within which a time- and/or space-varying fine structure constant can be readily admitted (Bekenstein, 1982; Carroll, 1998; Marciano, 1984; Brax et al., 2003) for many physicists the prior probability of such must be considered small, given both the results of previous experiments and our faith in certain long-standing physical principles. In particular, past analyses of the Oklo natural fission reactor (Shlyakhter, 1976; Damour and Dyson, 1996; Gould et al., 2006), Earth-fallen meteorite samples (Olive et al., 2004), and optical atomic clocks (Fortier et al., 2007; Rosenband et al., 2008) have strongly favored a scenario of negligible *temporal variation* in α locally; with Rosenband et al. (2008), for instance, determining $|\Delta\alpha/\alpha| < 1.6 \times 10^{-17}$ per year *at present on Earth*. These results may “daringly” (with respect to the strong assumption of zero higher derivatives) be extrapolated to $|\Delta\alpha/\alpha| < 2 \times 10^{-7}$ since the Big Bang (~ 13.4 Gyr ago); with more prosaic but reliable constraints of $|\Delta\alpha/\alpha| < 0.1$ over this time provided by contemporary analyses of fluctuations in the Cosmic Microwave Background (Rocha et al., 2004; Nakashima, Nagata and Yokoyama, 2008; Calabrese et al., 2011).

Spatial variation in the fine structure constant, on the other hand, has been previously constrained to $|\Delta\alpha/\alpha| < 10^{-4}$ on cosmic scales via both emission and absorption line based quasar studies (Bahcall and Salpeter, 1965; Bahcall, Sargent and Schmidt, 1967; Ivanchik, Potekhin and Varshalovich, 1999), although this may only be considered weak prior evidence against the Webb et al. team’s specific claim as the dipole strength inferred from the quasar dataset contributes an effect well below this level. More importantly from an astronomical perspective is our prior belief in the *cosmological principle*—namely, that the physical laws of the Universe are uniform throughout; and consequently that the Universe appears essentially homogeneous when viewed on sufficiently large scales. This principle not only serves as a key plinth of the predominant cosmological model but has been hitherto well-supported by measurements of the fractal dimension (converging toward three) at large scales in galaxy redshift surveys (Martínez et al., 1998; Scrimgeour et al., 2012) and by the isotropy of the Cosmic Microwave Background (Mandolesi et al. 1986; though see Eriksen et al. 2007 and Clarkson and Barrett 1999). The proposal of a large-scale spatial variation across the observable Universe in one of the fundamental coupling constants of QED is thus in direct conflict with this trusted paradigm. In conjunction with Sean Carroll’s previously-noted objections (from his “Cosmic Variance” blog; see Introduction) regarding the surprisingly low mass-energy budget implied by such a smoothly varying scalar field we must from this perspective assume

a significant prior weighting *against* the dipole interpretation.

To quantify this prior weighting we might for arguments sake suppose a preference approaching the 5σ level for the established theory inspired by the agreed evidence benchmark in the particle physics community for a “definite discovery” of the Higgs boson¹¹—that is, $P_{\text{prior}}(\text{strict null})/P_{\text{prior}}(\text{dipole}) \approx 3.5 \times 10^6$ (from the one-sided Normal tail). In this conservative scenario, even the $\sim 4\sigma$ support claimed by the Webb et al. team under their nominal error model (and broadly “confirmed” in this study with a Bayes factor of ≈ 180 , “corresponding” to $\approx 2.5\sigma$ support) must be considered insufficient to conclude that the fine structure constant varies across the Universe in the manner described. Indeed, such a prior weighting against the dipole hypothesis appears reflected in the relative caution with which both independent cosmologists (Olive, Peloso and Uzan, 2011) and the Webb et al. team themselves have at times discussed their results. Dedicated observations targeting quasars *at the poles of the inferred dipole* have thus been proposed (Webb et al., 2011; King et al., 2012) in order to establish the further support necessary for this hypothesis to become widely accepted.

However, as we have already demonstrated in Section 4.3 above the present quasar dataset does not exclude (in fact it weakly favors) the skeptical interpretation that there exist sources of opposing systematic bias in the $\Delta\alpha/\alpha$ estimates deriving from the Keck and VLT telescopes, artificially inducing the apparent dipole signature. And, crucially, such future observations¹² focussed exclusively on the poles can do little to settle this debate, even if the nominal error model and dipole hypothesis are in fact correct. We demonstrate this explicitly via the following simulation procedure making use of our posterior predictive, the results of which are shown in Figure 10. First we draw a vector of θ_m parameters for the dipole hypothesis and θ_e parameters for error model $\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$ from the (present) posterior. Using these as input we simulate n VLT data values ($\Delta\alpha/\alpha$ estimates) under this nominal error model for sightings along the maximum pole, and we assign each a z_{abs} and σ_{obs} sampled (with replacement) from the observed distribution in the quasar dataset; and we do the same thereafter for n replicate Keck (LC) observations sighted along the minimum pole. We then add all these mock observations to the existing quasar dataset, and recompute marginal likelihoods for the dipole hypothesis under $\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$ and the strict null under

¹¹Acknowledging that the 5σ significance benchmark of the particle physics community is in fact in reference to a long-run sampling p -value under the null.

¹²We assume these to be conducted on the same telescopes as the original dataset—namely, the Keck and the VLT—as these represent the most powerful (and efficient) facilities available at present for this type of spectroscopy.

AAA (using the very fast nested importance sampling algorithm). By repeating these steps a thousand times over we thus estimate the distribution of such future (log) Bayes factors¹³ under the proposed observing strategy (and error model–hypothesis pairing). As is evident from the lefthand panel of Figure 10, with even 250 new observations at each pole there remains less than a 15% chance of recovering more than 5σ Bayes factor support for the dipole with nominal error model (assumed true) over the strict null with skeptical error model. [We note that the wide range of predicted Bayes factors recovered here arises primarily from the present range of permitted dipole strengths expressed by our posterior on parameter B (see Figure 8), which dominates over the (well known) sampling variance of the Bayes factor itself (cf. Good 1992; Jenkins and Peacock 2011).]




FIG 10. The predicted power of future quasar observations for dismissing the skeptical interpretation (of opposing biases in the Keck and VLT error terms) supposing the dipole hypothesis with nominal (unbiased) error model as truth. In the lefthand panel we illustrate the simulated distribution of log marginal likelihood differences (i.e., log Bayes factors) recovered after the addition of 50, 100, and 250 additional mock observations ($\Delta\alpha/\alpha$ estimates) from each telescope directed along the corresponding pole given a random parameterization of the monopole+ $r(z)$ -dipole hypothesis drawn from its present posterior; each subject to the appropriate noise term under model $\mathbf{A}_0\mathbf{A}_0\mathbf{A}_0$ and sampled in z_{abs} and σ_{obs} as per the original dataset. We present the same in the righthand panel, but for a scenario in which identical numbers of total mock observations are split over both polar and equatorial sightings. The latter strategy clearly offers far better prospects for dismissing the skeptical hypothesis (if indeed false). For reference, the red, yellow, and green-shaded zones under each curve mark the $\Delta \log Z$ limits corresponding to $>3\sigma$, $>4\sigma$, and $>5\sigma$ support, respectively.

A more profitable strategy for distinguishing between the dipole under the nominal error model and the null under the default skeptical error model

¹³Trotta (2007) has proposed for this the term, “predicted posterior odds distribution”, with acronym, PPOD.

would instead be to focus in part on adding further observations along the equatorial strip where the relative bias between these two instruments may be most readily constrained. We once again demonstrate this by repeating our simulation procedure to predict future Bayes factors, but this time employing an observational strategy in which half the quasar sightings are placed on the equator (and half again on the poles). Our results are summarized in the righthand panel of Figure 10 from which it is evident that we now have a better than one-in-two chance of recovering the sought after 5σ support for the nominal dipole hypothesis (assumed as truth) for the $n = 250$ experiment.

Thus, we reach the final conclusions of our Bayesian reanalysis of the quasar dataset. Namely, that: (I) *given both our incomplete understanding of the observational errors and our limited theoretical (prior) expectations regarding the properties of any spatial variation in the fine structure constant the present observational coverage (featuring limited overlap of the two telescopes for which opposing biases might be suspected) must be deemed inadequate to properly distinguish the Webb et al. team's proposed dipole field from the (strict or monopole) null*; and (II) *one cannot afford to overlook the importance of observations along the equator of the alleged dipole in addition to those proposed along its poles when planning future campaigns (with the Keck and VLT telescopes) for the purpose of settling this debate.*

We have also demonstrated in this study the utility of the nested importance sampling technique for marginal likelihood computation: in particular with regard to (I) the ease with which it allows for testing the sensitivity of the Bayes factor to the priors; and (II) its potential to facilitate prediction of the Bayes factor distribution for future observations (under the specification of a particular hypothesis as truth) without unreasonable demand on computational power or time.

References.

- ALBRECHT, A. and MAGUELJO, J. (1999). Time varying speed of light as a solution to cosmological puzzles. *Physical Review D* **59** 043516.
- ANDRAE, R., SHULZE-HARTUNG, T. and MELCHIOR, P. (2010). Dos and don'ts of reduced chi-squared. Preprint(arXiv:1012.3754).
- BAHCALL, J.N. and SALPETER, E.E. (1965). On the interaction of radiation from distant sources on the intervening medium. *The Astrophysical Journal* **142** 1677–1680.
- BAHCALL, J.N., SARGENT, W.L.W. and SCHMIDT, M. (1967). An analysis of the absorption spectrum of 3C 191. *The Astrophysical Journal* **149** L11–L15.
- BARROW, J.D. (1999). Cosmologies with varying light speed. *Physical Review D* **59** 043515.
- BARROW, J.D. and MAGUELJO, J. (2000). Can a changing α explain the supernovae results? *The Astrophysical Journal* **532** L87–L90.
- BEKENSTEIN, J.D. (1982). Fine-structure constant: Is it really a constant? *Physical Review D* **25** 1527–1539.

- BERENGUT, J.C., KAVA, E.M. and FLAMBAUM, V.V. (2012). Is there a spatial gradient in values of the fine-structure constant? A reanalysis of the results. Preprint(arXiv:1203.5891).
- BORIE, E. (1986). The role of the fine-structure constant in QED. *Metrologia* **22** 140–145.
- BOSCH, J.F. (2011). Modeling techniques for measuring galaxy properties in multi-epoch surveys. Preprint(arXiv:1109.1033).
- BOUCHENDIRA, R., CLADÉ, P., GUELLATI-KHÉLIFA, S., NEZ, F. and BIRABEN, F. (2011). New determination of the fine structure constant and test of the quantum electrodynamics. *Physical Review Letters* **106** 080801.
- BRAX, Ph., VAN DE BRUCK, C., DAVIS, A.-C. and RHODES, C.S. (2012). Varying constants in brane world scenarios. *Astrophysics and Space Science* **283** 627–632.
- BURBIDGE, E.M., LYNDS, C.R. and BURBIDGE, G.R. (1966). On the measurement and interpretation of absorption features in the spectrum of the quasi-stellar object 3C 191. *The Astrophysical Journal* **144** 447–450.
- CARROLL, S.M. (1998). Quintessence and the rest of the world: Suppressing long-range interactions. *Physical Review Letters* **81** 3067–3070.
- CALABRESE, E., MENEGONI, E., MARTINS, C.J.A.P., MELCHIORRI, A. and ROCHA, G. (2011). Constraining variations in the fine structure constant in the presence of early dark energy. *Physical Review D* **84** 023518.
- CARLIN, B.P. and CHIB, S. (1995). Bayesian model choice via Markov Chain Monte Carlo Methods. *Journal of the Royal Statistical Society B* **57** 473–484.
- CARLIP, S. and VAIDYA, S. (2003). Cosmology: Do black holes constrain varying constants? *Nature* **421** 498–499.
- CHIPMAN, H., GEORGE, E.I. and MCCULLOCH, R.E. 2001. Practical implementation of Bayesian model selection. *IMS Lecture Notes – Monograph Series* **38** 67–134.
- CHOPIN, N. and ROBERT, C.P. (2010). Properties of nested sampling. *Biometrika* **97** 741–755.
- CLARKSON, C.A. and BARRETT, R.K. (1999). Does the isotropy of the CMB imply a homogeneous universe? Some generalized EGS theorems. *Classical and Quantum Gravity* **16** 3781–3794.
- DAMOUR, T. and DYSON, F. (1996). The Oklo bound on the time variation of the fine-structure constant revisited. *Nuclear Physics B* **480** 37–54.
- DAVIES, P.C.W., DAVIS, T.M. and LINEWEAVER, C.H. (2002). Cosmology: Black holes constrain varying constants. *Nature* **418** 602–603.
- DZUBA, V.A., FLAMBAUM, V.V. and WEBB, J.K. 1999. Calculations of the relativistic effects in many-electron atoms and space-time variation of fundamental constants. *Physical Review A* **59** 230–237.
- ERIKSEN, H.K., BANDAY, A.J., GÓRSKI, K.M., HANSEN, F.K. and LILJE, P.B. (2007). Hemispherical power asymmetry in the third-year Wilkinson Microwave Anisotropy Probe sky maps. *The Astrophysical Journal Letters* **660** 81–84.
- FEROZ, F. and HOBSON, M.P. (2007). Multimodal nested sampling: an efficient and robust alternative to Markov Chain Monte Carlo methods for astronomical data analyses. *Monthly Notices of the Royal Astronomical Society* **384** 449–463.
- FEROZ, F., HOBSON, M.P. and BRIDGES, M. (2009). MULTINEST: an efficient and robust Bayesian inference tool for cosmology and particle physics. *Monthly Notices of the Royal Astronomical Society* **398** 1601–1614.
- FORTIER, T.M., ASHBY, N., BERGQUIST, J.C., DELANEY, M.J., DIDDAMS, S.A., HEAVNER, T.P., HOLLBERG, L., ITANO, W.M., JEFFERTS, S.R., KIM, K., LEVI, F., LORINI, L., OSKAY, W.H., PARKER, T.E., SHIRLEY, J. and STALKNER, J.E. (2007). Precision atomic spectroscopy for improved limits of variation of the fine structure constant and local

- position invariance. *Physical Review Letters* **98** 070801.
- FRIEL, N. and PETTITT, A.N. (2008). Marginal likelihood estimation via power posteriors. *Journal of the Royal Statistical Society B* **70** 589–607.
- FRIEL, N. and WYSE, J. (2012). Estimating the evidence – a review. *Statistica Neerlandica* ??? ??–??.
- GELMAN, A. and MENG, X.-L. (1998). Simulating normalizing constants: from importance sampling to bridge sampling to path sampling. *Statistical Science* **13** 163–185.
- GOOD, I.J. (1992). The Bayes/non-Bayes compromise: A brief review. *Journal of the American Statistical Association* **87** 597–606.
- GOULD, C.R., SHARAPOV and E.I., LAMOREAUX, S.K. (2006). Time variability of α from realistic models of Oklo reactors. *Physical Review C* **74** 024607.
- GRIEST, K., WHITMORE, J.B., WOLFE, A.M., PROCHASKA, J.X., HOWK, J.C. and MARCY, G.W. (2010). Wavelength accuracy of the Keck HIRES spectrograph and measuring changes in the fine structure constant. *The Astrophysical Journal* **708** 158–170.
- GRIFFITHS, D.J. (2005). *Introduction to Quantum Mechanics*, 2nd ed. Pearson Prentice Hall, Upper Saddle River, NJ.
- HANNEKE, D., FOGWELL, S. and GABRIELSE, G. (2008). New measurement of the electron magnetic moment and the fine structure constant. *Physical Review Letters* **100** 120801.
- HOGG, D.W. (1999). Distance measures in cosmology. Preprint(arXiv:9905116).
- IVANCHIK, A.V., POTEKHIN, A.Y. and VARSHALOVICH, D.A. (1999). The fine-structure constant: a new observational limit on its cosmological variation and some theoretical consequences. *Astronomy and Astrophysics* **343** 439–445.
- JAYNES, E.T. (2003). Probability theory: The logic of science. *Cambridge University Press*, Cambridge, UK.
- JEFFREYS, W.H. and BERGER, J.O. (1992). Ockham’s razor and Bayesian analysis. *American Scientist* **80** 64–72.
- JENKINS, C.R. and PEACOCK, J.A. (2011). The power of Bayesian evidence in astronomy. *Monthly Notices of the Royal Astronomical Society* **413** 2895–2905.
- KASS, R.E. and RAFTERY, A.E. (1995). Bayes Factors. *Journal of the American Statistical Association* **90** 773–795.
- KENDALL, D. (1938). Effect of radiation damping on doppler broadening on the atomic absorption coefficient. *Zeitschrift für Astrophysik* **16** 308–317.
- KING, J.A., MORTLOCK, D., WEBB, J. and MURPHY, M. (2010). Markov Chain Monte Carlo methods applied to measuring the fine structure constant from quasar spectroscopy. *Highlights of Astronomy* **15** 318–318.
- KING, J.A., WEBB, J.K., MURPHY, M.T., FLAMBAUM, V.V., CARSWELL, R.F., BAINBRIDGE, M.B., WILCZYNSKA, M.R., and KOCH, F.E. (2012). Spatial variation in the fine structure constant – new results from VLT/UVES. *Monthly Notices of the Royal Astronomical Society* **422** 3370–3414.
- LAMOREAUX, S.K. and TORGERSON, J.R. (2004). Neutron moderation in the Oklo natural reactor and the time variation of alpha. *Physical Review D* **69** 121701.
- LEFEBVRE, G., STEELE, R. and VANDAL, A.C. (2010). A path sampling identity for computing the Kullback-Leibler and J divergences. *Computational Statistics & Data Analysis* **54** 1719–1731.
- LEVSHAKOV, S.A., COMBES, F., BOONE, F., AGAFONOVA, I.I., REIMERS, D. and KOZLOV, M.G. (2012). An upper limit to the variation in the fundamental constants at redshift $z = 5.2$. *Astronomy & Astrophysics* **540** L9.
- MALLOWS, C.L. (1995). More comments on C_P . *Technometrics* **37** 362–372.
- MANDOLESI, N., CALZOLARI, P., CORTIGLIONI, S., DELPINO, F., SIRONI, G., INZANI, P., DEAMICI, G. and 6 coauthors (1986). Large-scale homogeneity of the Universe measured

- by the microwave background. *Nature* **319** 751–753.
- MARCIANO, W. (1984). Time variation of the fundamental “constants” and Kaluza-Klein theories. *Physical Review Letters* **52** 489–491.
- MARTÍNEZ, V.J., PONS-BORDERÍA, M.-J., MOYED, R.A. and GRAHAM, M.J. (1998). Searching for the scale of homogeneity. *Monthly Notices of the Royal Astronomical Society* **298** 1212–1222.
- MOFFAT, J.W. (1993). Superluminary universe: A possible solution to the initial value problem in cosmology. *International Journal of Modern Physics D* **2** 351–365.
- MOLARO, P., LEVSHAKOV, S.A., MONAI, S., CENTURIÓN, M., BONIFACIO, P., D’ODORICO, S. and MONACO, L. (2008). UVES radial velocity accuracy from asteroid observations. *Astronomy and Astrophysics* **481** 559–569.
- MUKHERJEE, P., PARKINSON, D. and LIDDLE, A.R. (2006). A nested sampling algorithm for cosmological model selection. *The Astrophysical Journal Letters* **638** 51–54.
- MURPHY, M.T., WEBB, J.K., FLAMBAUM, V.V., PROCHASKA, J.X. and WOLFE, A.M. (2001). Further constraints on variation of the fine-structure constant from alkali-doublet QSO absorption lines. *Monthly Notices of the Royal Astronomical Society* **327** 1237–1243.
- MURPHY, M.T., WEBB, J.K. and FLAMBAUM, V.V. (2003). Further evidence for a variable fine-structure constant from Keck/HIRES QSO absorption spectra. *Monthly Notices of the Royal Astronomical Society* **345** 609–638.
- NAKASHIMA, M., NAGATA, R. and YOKOYAMA, J. (2008). Constraints on the time variation of the fine structure constant by the 5-year WMAP data. *Progress of Theoretical Physics* **120** 1207–1215.
- NASON, G.P. (2006). On the sum of t and Gaussian random variables. *Statistics and Probability Letters* **76** 1280–1286.
- OLIVE, K.A., POSPELOV, M., QIAN, Y.-Z., MANHÈS, G., VANGIONI-FLAM, E., COC, A. and CASSÉ, M. (2004). Reexamination of the ^{187}Re bound on the variation of fundamental couplings. *Physical Review D* **69** 027701.
- OLIVE, K.A., PELOSO, M. and UZAN, J.-P. (2011). A wall of fundamental constants. *Physical Review D* **83** 43509.
- PEWSEY, A. (2000). The wrapped skew-normal distribution on the circle. *Communications in Statistics – Theory and Methods* **29** 2459–2472.
- ROCHA, G., TROTTA, R., MARTINS, C.J.A.P., MELCHIORRI, A., AVELINO, P.P., BEAN, R. and VIANA, P.T.P. (2004). Measuring α in the early universe: cosmic microwave background polarization, re-ionization and the Fisher matrix analysis. *Monthly Notices of the Royal Astronomical Society* **352** 20–38.
- ROSENBAND, T., HUME, D.B., SCHMIDT, P.O., CHOU, C.W., BRUSCH, A., LORINI, L., OSKAY, W.H., BRULLINGER, R.E., FORTIER, T.M., STALNAKER, J.E., DIDDAMS, S.A., SWANN, W.C., NEWBURY, N.R., ITANO, W.M., WINELAND, D.J. and BERGQUIST, J.C. (2008). Frequency ratio of Al^+ and Hg^+ single-ion optical clocks; Metrology at the 17th decimal place. *Science* **319** 1808–1812.
- ROUSSEUW, P.J. (1984). Least median of squares regression. *Journal of the American Statistical Association* **74** 871–880.
- SAVEDOFF, J.D. (1956). Physical constants in extra-galactic nebulae. *Nature* **178** 688–689.
- SCRIMGEOUR, M., DAVIS, T., BLAKE, C., JAMES, J.B., POOLE, G., STAVELEY-SMITH, L., BROUGH, S., COLLESS, M. and 21 coauthors (2012). The WiggleZ Dark Energy Survey: the transition to large-scale cosmic homogeneity. Preprint(arXiv:1205.6812).
- SHLYAKHTER, A.I. (1976). Direct test of the constancy of fundamental nuclear constants. *Nature* **264** 340.
- SKILLING, J. (2006). Nested sampling for general Bayesian computation. *Bayesian Analysis*

- 1** 833-860.
- STOCKTON, A.N. and LYND, C.R. (1966). The remarkable absorption spectrum of 3C 191. *The Astrophysical Journal* **144** 451–453.
- TROTTA, R. (2007). Forecasting the Bayes factor of a future observation. *Monthly Notices of the Royal Astronomical Society* **378** 819–824.
- WEBB, J.K., FLAMBAUM, V.V., CHURCHILL, C.W., DRINKWATER, M.J. and BARROW, J.D. (1999). Search for time variation of the fine structure constant. *Physical Review Letters* **82** 884–887.
- WEBB, J.K., MURPHY, M.T., FLAMBAUM, V.V., DZUBA, V.A., BARROW, J.D., CHURCHILL, C.W., PROCHASKA, J.X. and WOLFE, A.M. (2001). Further evidence for cosmological evolution of the fine structure constant. *Physical Review Letters* **87** 091301.
- WEBB, J.K., KING, J.A., MURPHY, M.T., FLAMBAUM, V.V., CARSWELL, R.F. and BAINBRIDGE, M.B. (2011). Indications of a spatial variation of the fine structure constant. *Physical Review Letters* **107** 191101.
- WEISS, A., WALTER, F., DOWNES, D., CARILLI, C.L., HENKEL, C., MENTEN, K.M. and COX, P. (2012). On the variations of fundamental constants and AGN feedback in the QSO host galaxy RXJ0911.4+0551 at $z = 2.79$. Preprint(arXiv:1204.5614).
- WHITMORE, J.B., GRIEST, K. and MURPHY, M.T. (2010). Wavelength calibration of the VLT-UVES spectrograph. *The Astrophysical Journal* **723** 89–99.
- WILCOXON, F. (1945). Individual comparisons by rank methods. *Biometrics Bulletin* **1** 80–83.

QUEENSLAND UNIVERSITY OF TECHNOLOGY
 SCHOOL OF MATHEMATICAL SCIENCES (STATISTICAL SCIENCE)
 GPO Box 2434, BRISBANE 4001
 QLD, AUSTRALIA
 E-MAIL: dr.ewan.cameron@gmail.com